

Research article

Acute Effects of Dynamic Stretching Followed by Vibration Foam Rolling on Sports Performance of Badminton Athletes

Wei-Cheng Lin¹, Chia-Lun Lee² and Nai-Jen Chang^{1,3,4}✉

¹ Department of Sports Medicine, Kaohsiung Medical University, Kaohsiung 807, Taiwan

² Center for Physical and Health Education, National Sun Yat-sen University, Kaohsiung 804, Taiwan

³ PhD Program in Biomedical Engineering, Kaohsiung Medical University, Kaohsiung 807, Taiwan

⁴ Regenerative Medicine and Cell Therapy Research Center, Kaohsiung Medical University, Kaohsiung 807, Taiwan

Abstract

Dynamic stretching (DS) is performed to increase sports performance and is also used primarily for transiently increasing range of motion (ROM). Recently, vibration foam rolling (VFR) has emerged. Its underlying concept is that it combines foam rolling techniques with local vibration to improve ROM and muscular activation concurrently. This crossover study investigated the effects of DS or DS followed by VFR (DS + VFR) during warm-ups on flexibility, muscle stiffness, power, and agility of the lower limbs in badminton athletes. Forty badminton players performed DS or DS + VFR as warm-up exercises on two occasions in a randomized order. The target muscle groups were the bilateral shoulder, anterior and posterior thigh, posterior calf, and lower back. Main outcome measures: The primary outcome was knee range of motion (ROM), and the secondary outcomes were muscle stiffness, lower limb power (countermovement jump [CMJ]), and agility. Results indicated that the protocols improved performance. DS increased knee flexion ROM (% change = 1.92, ES = 0.3, $p = 0.033$), CMJ height (% change = 5.04, ES = 0.2, $p = 0.004$), and agility (% change = -4.97, ES = 0.4, $p < 0.001$) but increased quadriceps muscle stiffness (% change = 3.74, ES = 0.3, $p = 0.001$) and increased gastrocnemius muscle stiffness (% change = 10.39, ES = 0.5, $p = 0.001$). DS + VFR increased knee extension ROM (% change = 2.87, ES = 0.4, $p = 0.003$), reduced quadriceps muscle stiffness (% change = -2.79, ES = 0.3, $p = 0.017$), CMJ height (% change = 2.41, ES = 0.1, $p = 0.037$), and agility (% change = -4.74, ES = 0.2, $p < 0.001$). DS + VFR was not significantly superior to DS, except for muscle stiffness reduction. Taken together, we suggest that practitioners consider DS as a first line of warm-up exercise to increase ROM, CMJ height, and agility in athletes. Moreover, the addition of VFR to DS results in a large reduction of muscle stiffness, potentially reducing the risk of sports injury. Athletes, coaches and athletic professionals may consider them when selecting effective warm-up practices to augment athletic performance.

Key words: Warm up exercise, foam rolling, athletic performance, sports, vibration therapy

Introduction

Badminton is a popular sport and one of the fastest racquet sports (Lees, 2003). The lunge step is one of the most frequently performed movements in badminton, accounting for approximately 15% of the total movements in a game (Kuntze et al., 2010; Mei et al., 2017). Performing a good lunge step is usually associated with high flexibility. In

addition, badminton players must react by moving rapidly with powerful jumps and agile footwork throughout a game. Players must repeat actions quickly with high speed and intensity. Therefore, players need excellent joint range of motion (ROM), power, and agility (Tiwari et al., 2011; Wong et al., 2019).

Warm-up exercise is critical prior to participation in sports. It improves performance and may avoid injury (Woods et al., 2007). Muscle strain injuries usually occur during movements that involve rapid acceleration/deceleration and sprinting (Opar et al., 2012). Dynamic stretching (DS) is the most commonly suggested warm-up protocol (Turki et al., 2019). The DS technique involves a stretch to lengthen the muscle, and it is performed by moving parts of the body and gradually increasing reach and speed of movement (Behm and Chaouachi, 2011). It often mimics movement patterns performed during subsequent exercise. DS provides a more sport-specific warm-up exercise, and as a precursor, it increases body temperature, improves nerve conduction, and increases sports performance (i.e., jump height, sprint speed, and agility) (Chaouachi et al., 2010; Opplert and Babault, 2018; Perrier et al., 2011). However, reports regarding the effect of DS on muscle stiffness are conflicting, indicating that DS may cause increased or reduced muscle stiffness (Chen et al., 2018; Iwata et al., 2019; Pamboris et al., 2018).

Vibration foam rolling (VFR) combines a foam roller with vibration; the roller serves as a foam rolling tool that targets a particular muscle group. It has been suggested as an alternative warm-up method (Lee et al., 2018; Lyu et al., 2020). Beneficial outcomes of VFR include remobilizing soft-tissue compliance to enable longer muscle length and increasing blood flow and circulation to soft tissues conducting rolling on the soft tissue (MacDonald et al., 2013). Concurrently, the additional transmission of mechanical oscillations to the target muscle may increase the number of motor units recruited (Cochrane, 2011). However, few studies have investigated the effectiveness of VFR, particularly for athletic performance. Thus far, studies have focused on ROM (Cheatham et al., 2019; Garcia-Gutierrez et al., 2018; Lee et al., 2018; Lim et al., 2019; Lim and Park, 2019; Lyu et al., 2020), pressure pain threshold (Cheatham et al., 2019; Romero-Moraleda et al., 2019), electromyography activity (Lim et al., 2019), isokinetic muscle strength (Lee et al., 2018; Lyu et al., 2020), perceived joint stability (De Benito et al., 2019), dynamic balance (Lee et al., 2018), and vertical jump (Lim and Park,

2019). However, these studies have focused on the knee or ankle musculatures rather than combined effects on the whole body—specifically, when the technique targets upper and lower extremities and the lower back. Furthermore, the aforementioned VFR studies were conducted in healthy participants or participants involved in recreational activities rather than in athletes. Scientific evidence is inadequate; little is known about sports performance following the use of VFR as a warm-up regime.

The most beneficial warm-up protocol remains uncertain, particularly for combination warm-up exercises. Moreover, no study has examined the effects of DS combined with VFR on badminton athletes. Studying the effects of combination warm-ups on sports performance can provide evidence that can help athletes, coaches, and clinical professionals to decide whether to add the VFR protocol to DS. Accordingly, this study compared the acute effects of DS as well as DS followed by VFR during warm-up on flexibility, muscle stiffness, power, and agility in young adults. The primary outcome was knee ROM. The secondary outcomes were muscle stiffness, countermovement jump (CMJ), and agility. We hypothesized that DS as warm-up exercise can not only increase ROM, CMJ height, and agility, but increase muscle stiffness. In addition, VFR can offset muscle stiffness.

Methods

Participants

The study protocol was approved by the Yuan's General Hospital Institutional Review Board (Approval Number: 20180209B) and Human Research Ethics Committee in the spirit of the Helsinki Declaration. Participants were informed of the benefits and risks of the study, and they signed an informed consent form before participating. Totally, 40 college badminton players (25 male and 15 female students) participated in two trials: (1) DS and (2) DS followed by VFR (Figure 1). Characteristics of participants are presented in Table 1. The inclusion criteria were as follows: membership in a school badminton team, 20–30 years of age, and no incidence of musculoskeletal disorder in the preceding 6 months. The exclusion criteria were as follows: musculoskeletal disease or neurological impairment, cardiovascular or respiratory disease, irradiation pain/radiation pain transferred to the lower extremities, previous surgery, and taking medication (e.g., anti-inflammatory or muscle relaxants) during the preceding 6 months.

Study procedures

This study was a crossover study with a within-subject design. Tests were performed in an indoor badminton gym at National Sun Yat-sen University. Each participant did the exercises at approximately the same time of day (18:00–20:00). Before the assessment session, participants performed a familiarization session, in which they were instructed by a certified physiotherapist on how to perform DS and VFR exercise regimes. During this orientation, participants were familiarized with the procedures and practiced with the assessment tools and equipment. Participants

were asked to maintain normal training programs but to avoid vigorous exercise at least 24 hours before the tests. Each participant performed the two protocols on separate occasions in a randomized order, with an interval of 48 hours (Su et al., 2017). Participants were individually guided by a certified physical therapist for performing both DS and VFR. The target muscle groups were the bilateral shoulder, anterior thigh, posterior thigh, posterior calf, and lower back. Participants performed the pretests in the following order: muscle stiffness, flexibility, CMJ, and agility tests. After completion of pretests, participants did the warm-up exercise (DS or DS + VFR) in a counterbalanced order. Posttest measurements were conducted in the same order as pretest measurements.

Table 1. Characteristics of participants Values are mean (standard deviation)

	Total (n = 40)	Male (n = 25)	Female (n = 15)
Age (year)	21.4 (1.5)	21.2 (1.5)	21.6 (1.6)
Height (m)	1.70 (0.10)	1.74 (0.07)	1.62 (0.07)
Body weight (kg)	64.8 (10.6)	69.5 (9.8)	56.8 (6.2)
BMI (kg/m ²)	22.4 (2.5)	22.8 (2.7)	21.8 (1.9)
Badminton experience (y)	7.1 (3.4)	7.1 (3.3)	7.1 (3.8)

Outcome measures

Before data collection, examiners were well-trained in conducting the flexibility, muscle stiffness, CMJ, and agility tests.

Primary outcome

The primary outcome was knee ROM. The knee flexion ROM of participants was measured using Ely's test, which is also used for testing quadriceps femoris muscle flexibility. This test was shown to have an intraclass correlation coefficient (ICC) of 0.91, which represents high reliability (Piva et al., 2006). With participants in the prone position, the examiner aligned the plastic goniometer with the axis of the knee joint; the stationary arm represents the greater trochanter of the femur, and the passively moving arm represents the lateral malleolus of the ankle. When the hip flexes as the knee is flexed, the pelvis should remain on the floor, and compensation by hyper-lordosis of the lumbar spine is avoided. The outcome measured was rectus femoris tightness by the same researcher, which was used to determine the termination of knee flexion ROM. The researcher subjectively perceived to inability to push the leg farther without any body compensation and measured the ROM. The average measurement of two trials was recorded. In this study, the ICC was 0.821, suggesting high test–retest reliability. In addition, the minimum detectable change (MDC) value was calculated to be 2.2°, corresponding to a standard error of measurement (SEM) of 0.989.

The knee extension ROM of participants was measured using the popliteus angle test, which is also used to measure hamstring flexibility. This test was shown to have an ICC of 0.90, representing high reliability (Youdas et al., 2005). With participants in the supine position, the examiner aligned the plastic goniometer with the axis of

the knee joint. The stationary arm represents the vertical ground, and the moving arm represents the lateral malleolus of the ankle. The knee was actively moved upward until the participant felt some tightness but not pain, and the angle between the thigh and calf was measured. The average measurement of two trials was recorded. The ICC was 0.99, suggesting excellent test–retest reliability. The MDC value was calculated to be 2.334°, corresponding to an SEM of 0.842.

Secondary outcomes

Secondary outcomes included muscle stiffness test, CMJ test, and agility test.

Regarding muscle stiffness test, we used a handheld myometer (Myoton® PRO; Myoton AS, Tallinn, Estonia) to measure muscle stiffness. The myometer was shown to have high to excellent reliability (Aird et al., 2012). Myometer measurements were taken by holding the device above the skin overlaying the target muscle assessment site. Once the desired position was achieved, mechanical impact (duration: 15 milliseconds; force: 0.4 N) was delivered to the muscle using a mechanical probe, causing the tissue to briefly deform (Pruyn et al., 2016). The target muscle groups included the middle point of the quadriceps and gastrocnemius. Stiffness was calculated as the ratio between the force applied and muscle deformation (Ditroilo et al., 2011). The average measurement of two trials was recorded. The gastrocnemius ICC was 0.910, and the quadriceps ICC was 0.983, suggesting good to excellent test–retest reliability.

Regarding CMJ test, My jump is a software application available on Appstore (Apple Inc., USA), and it is used to measure CMJ height through frame-by-frame analysis. It has excellent reliability (ICC = 0.997) (Balsalobre-Fernández et al., 2015). The in-app settings allow slow-motion playback for easy identification of the video frames in which jump take-off and landing occur (Stanton et al., 2015). Participants made vertical jumps, with feet shoulder-width apart and toes aligned behind a white line. Participants placed their hands on the hips and kept their knees straight during flight while jumping as high as possible.

Three jumps were recorded, with a resting period of 1 minute. The highest height was selected for analysis. The ICC was 0.959, suggesting excellent test–retest reliability. Regarding agility test, FITLIGHT Trainer™ (FITLIGHT Sports Corp., Aurora, ON, Canada) was adopted to measure agility (McMillian et al., 2006a). It is a wireless reaction training system consisting of LED-powered lights controlled by a tablet. The lights can be deactivated by touch. During the test, participants used their dominant hand to touch a red light on the floor until the light deactivated. The test area was the size of a badminton half-court, with eight lights placed at corners and half lines (Figure 1). After touching a light, the player returned to the center of the area. Participants extinguished the eight lights in order as quickly as possible. The completion time was recorded. The ICC was 0.969, suggesting excellent test–retest reliability.

DS exercises

The DS protocol comprised eight movements through the active ROM of the trunk and upper and lower extremities, involved moving parts of the body, and gradually increasing reach, speed of movement, or both. The exercises are detailed in Figure 2. They consisted of controlled leg and arm swings to the limits of the participant's ROM. The exercises were performed from one sideline to the opposite sideline of a badminton court and back (total distance: 18 m), with 10-second intervals (Chatzopoulos et al., 2014).

VFR exercises

Participants were individually instructed by a physical therapist to use a vibrating roller (dimensions: 36 × 20 × 15 cm³; weight: 1.8 kg) that included a vibration generating motor (frequency: 28 Hz) enclosed by a polypropylene foam outer shell (Vyper, Hyperice, Irvine, CA, USA). Participants were asked to put as much body mass as was tolerable on the roller. They performed active and smooth rolling back and forth on the target muscle groups (Figure 2). Each muscle group received 20 seconds of vibration. The duration of single roll was 2 seconds. The exercises (except for that for the lower back) were executed on the opposite side (Figure 3).




Figure 1. Experimental court for the badminton agility test. The numbers ① to ⑧ represent light placement.









			
Arm crossover	Rotator cuff	Walking lunge with trunk rotation	Lateral shuffle
Stand with feet shoulder-width apart and swing arms bilaterally across the chest while moving forward.	Elevate the arm overhead with an elbow flexion and then gently lean to the side.	Perform a large step forward while concurrently rotating arms horizontally.	Move laterally without crossing feet.
			
Frankenstein walk	Heel-ups	Inch worms (hand walk)	Modified shuttle run
Extend hands in front of the body with palms down and kick toward the hands with legs extended.	Kick heels toward the buttocks while moving forward.	Start in push-up position. Keep legs extended and walk toward the hands then move forward while keeping the limbs stretched (repeated six times).	Run to the opposite line at a moderate speed (maximum speed of 50%), bend to touch the line, and then return with gradual acceleration (75%) and touch the starting line. After touching the starting line, run to the opposite line at near maximum speed (90%), touch the line, and walk to return to the starting line.

Figure 2. Dynamic stretching protocols.

Statistical Analyses

An a priori sample size calculation based on anticipated differences in knee extension ROM as the primary outcome was conducted using an anticipated difference of 4° with a standard deviation of 6° between pretest and postintervention in the DS + VFR group. The calculation, with an alpha level of 0.01 and a desired statistical power of 80%, was conducted using G*Power (Faul et al., 2007). The minimum sample size was 30. To increase statistical power, we enrolled 40 participants.

Statistical analyses were performed using SPSS version 20.0 (Chicago, IL, USA). Data are expressed as mean and standard deviation and tested for normality (Shapiro–Wilk’s test, $p > 0.05$), and homogeneity of variance was confirmed using Levene’s test. If results of the Mauchly’s sphericity test indicated the sphericity assumption was violated, the Greenhouse–Geisser adjustment was adopted to correct the degrees of freedom. A 2 (condition: DS vs. DS + VFR) × 2 (time: pretest vs. posttest) repeated

measures analysis of variance was conducted to test the effects of different conditions on dependent variables. If a significant group × time interaction was identified, follow-up analysis was conducted using a t test to determine the effect. Paired t tests within each group were conducted to determine significant main effects of the intervention. Furthermore, changes in values from pretest to posttest were calculated and analyzed using a paired t test. The effect size (Cohen’s d) was calculated to show the magnitude of the effects ($d = M1 - M2 / \sigma_{pooled}$) for each group. Significance level (α) was set at 0.05.

To confirm that ROM measurement was greater than measurement error, the MDC value was calculated as follows: SEM was initially calculated using the formula $SEM = SD \times \sqrt{1 - ICC}$, where SD is the standard deviation of scores from the first test, and ICC is the test–retest ICC. Subsequently, MDC was calculated using the following formula: $1.96 \times \sqrt{2} \times SEM$ (Chen et al., 2019).




Figure 3. Vibration rolling protocols.

Table 2. Outcomes for dynamic stretching with or without vibration foam rolling

Parameters	DS							DS + VFR					
	Pre	Post	Change	% Change	Effect size	p	Pre	Post	Change	% Change	Effect size	p	
Knee ROM	Flexion (degrees)	124.1 (7.2)	126.4 (7.4)*	2.38 (6.8)	1.92	0.3	0.033	122.8 (8.1)	123.2 (10.3)	0.4 (6.9)	0.33	0.1	0.692
	Extension (degrees)	143.0 (10.7)	146.8 (12.5)*	3.74 (9.7)	2.62	0.3	0.019	142.7 (9.4)	146.8 (10.5)*	4.1 (8.0)	2.87	0.4	0.003
Muscle stiffness	Quadriceps	258.9 (27.6)	268.5 (33.7)*	9.7 (16.5)	3.74	0.3	0.001	265.6 (23.7)	258.2 (32.1)*	-7.4 (18.7) [†]	-2.79	0.3	0.017
	Gastrocnemius	329.3 (61.7)	363.5 (70.2)*	34.2 (59.2)	10.39	0.5	0.001	314.6 (39.8)	314.6 (33.9)	0.1 (36.8) [†]	0.03	0	0.993
CMJ	Height (cm)	37.7 (9.5)	39.6 (10.5)*	1.9 (3.9)	5.04	0.2	0.004	37.4 (9.3)	38.2 (9.6)*	0.9 (2.5)	2.41	0.1	0.037
Agility	Total time (s)	18.1 (1.7)	17.3 (1.9)*	-0.9 (1.0)	-4.97	0.4	<0.001	19.0 (1.8)	18.7 (0.6)*	-0.9 (0.7)	-4.74	0.2	<0.001

* Significant difference ($p < 0.05$) compared with the pretest results. [†]Significant difference ($p < 0.05$) compared with DS.

Results

All outcomes are displayed in Table 2.

Knee ROM outcomes

For the knee flexion ROM, condition \times time interactions ($p = 0.25$) were not significant. However, significant main effects were observed for condition ($p = 0.002$) and time ($p = 0.051$). Compared with the pretest results, according to the post hoc test results, significant improvement in Ely's test was found in the DS group (% change = 1.92, ES

= 0.3, $p = 0.033$), but no significant improvement was identified in the DS+VFR group (% change = 0.33, ES = 0.1, $p = 0.629$).

For the knee extension ROM test, the condition \times time interaction ($p = 0.88$) and main effect of condition ($p = 0.89$) were not significant. However, a significant main effect of time ($p < 0.001$) was observed, indicating a difference between pre- and posttest knee extension ROM. Compared with the pretest results, according to the post hoc test results, both the DS group (% change = 2.62, ES = 0.3, $p = 0.019$) and DS + VFR group (% change = 2.87, ES

= 0.3, $p = 0.003$) exhibited significant improvement in the popliteus angle test.

Muscle stiffness outcomes

For quadriceps muscle stiffness, the condition \times time interaction was significant. The DS group demonstrated significantly increased stiffness compared with the DS + VFR group ($p < 0.001$). We further analyzed the simple main effect. Compared with the pretest results, the DS group exhibited a significant increase in quadriceps stiffness (% change = 3.74, ES = 0.3, $p = 0.001$), but the DS + VFR group showed decreased stiffness (% change = -2.79, ES = 0.3, $p = 0.017$).

For gastrocnemius stiffness, the condition \times time interaction was significant. The DS group demonstrated significantly increased stiffness compared with the DS + VFR group ($p = 0.001$). We analyzed the simple main effect. Compared with the pretest results, according to the post hoc test results, the DS group exhibited a significant increase in gastrocnemius stiffness (% change = 10.39, ES = 0.5, $p = 0.001$), but the DS + VFR group did not (% change = 0.03, ES = 0, $p = 0.993$).

CMJ outcomes

For the CMJ test, the condition \times time interaction ($p = 0.192$) and the main effect of condition ($p = 0.089$) were not significant. However, the main effect of time was significant ($p < 0.001$). Compared with the pretest results, according to the post hoc test results, participants exhibited significant improvement in CMJ height after the DS (% change = 5.04, ES = 0.2, $p = 0.004$) and DS + VFR (% change = 2.41, ES = 0.1, $p = 0.037$) interventions.

Agility outcomes

For the agility test, the condition \times time interaction in total time ($p = 0.625$) was not significant. However, the main effect of the condition ($p < 0.001$) and the time effect were significant ($p < 0.001$). Compared with the pretest results, according to the post hoc test results, participants exhibited significant improvement in agility after the DS (% change = -4.97, ES = 0.4, $p < 0.001$) and DS + VFR (% change = -4.74, ES = 0.2, $p < 0.001$) interventions.

Discussion

This is the first study to investigate the acute effects of DS + VFR as a whole-body warm-up protocol including the upper and lower limbs and trunk muscles in badminton athletes. Most research on VFR has focused on single joints of the body, such as the knee (Cheatham et al., 2019; De Benito et al., 2019; Lee et al., 2018; Lim et al., 2019; Lim and Park, 2019; Romero-Moraleda et al., 2019) or ankle (Garcia-Gutierrez et al., 2018; Lyu et al., 2020). The findings indicated that DS with or without VFR significantly increased knee ROM, CMJ height, and agility; however, DS increased thigh and calf muscle stiffness. DS followed by VFR did not have synergistic effects.

DS integrates controlled movement through the active ROM of a joint and includes change-of-direction movements. The current results for DS confirmed positive

effects, consistent with previous findings regarding flexibility (Paradis et al., 2014; Ryan et al., 2014a; Su et al., 2017), jump height (Fletcher, 2010; Ryan et al., 2014a; Turki et al., 2011), and agility (Chaouachi et al., 2010; McMillian et al., 2006b). The improved sports performance may be because DS increases muscle temperature or causes rearrangement/slipping of collagen fibers to regain flexibility (Herda et al., 2013; Nordez et al., 2009). In addition, DS provides a preload stimulus before an actual activity as post-activation potentiation. This generates increased phosphorylation of myosin light chains and motor neuron excitability (Hodgson et al., 2005); thus, athletic performance can be augmented.

An unanticipated finding was that the effects of DS + VFR were not superior to those of DS alone on flexibility, lower limb power, or agility in the athletes. Possible reasons are as follows: (1) optimal protocols for VFR (i.e., frequency, duration, vibration setting, and target muscles) have not been established; (2) the selection of participants (students, recreationally active adults, and athletes) differed among studies; and (3) outcomes of laboratory measures (i.e., isokinetic dynamometer) may differ from those of practical field tests. In the present study in badminton athletes, for the practical use of VFR, we selected the following protocol: 28 Hz, 20 seconds \times 1 set on the bilateral rotator cuff, quadriceps, hamstrings, posterior gastrocnemius, and lower back muscles). Moreover, we used field assessments for power and agility. In a laboratory assessment, Lyu et al. indicated that the use of VFR (28 Hz, 30 seconds \times 3 sets on bilateral gastrocnemius muscles) in recreationally active adults significantly improved ankle plantar flexor peak torque (Lyu et al., 2020). Lee et al. determined that the use of VFR (28 Hz, 30 seconds \times 3 sets on bilateral quadriceps and hamstring muscles) significantly increased quadriceps isokinetic muscle strength (Lee et al., 2018). However, Garcia-Gutiérrez et al. demonstrated that use of VFR (49 Hz, 20 seconds \times 3 sets on bilateral gastrocnemius muscles) in undergraduate students did not significantly increase ankle muscle maximal voluntary isometric contraction. Sağıroğlu et al. found no significant difference in jump height after the use of VFR (38 Hz, 30 seconds \times 2 sets on hamstrings, quadriceps, gluteals, and gastrocnemius) in well-trained male soccer players. Lim et al. indicated that the use of VFR (38 Hz, 60 seconds \times 5 sets on bilateral hamstrings) in college students caused no significant improvement in vertical jump (Lim and Park, 2019). Future studies should investigate optimal warm-up protocols for different populations and validate whether the beneficial effects identified in laboratory assessments can be translated to field applications.

Muscle stiffness is one of the main components of athletic performance (Miyamoto et al., 2017). Research has indicated that performing DS protocols after repeated muscle contractions may impair performance due to muscle fatigue (Costa et al., 2014; Ryan et al., 2014b) and may cause increased muscle stiffness (Pamboris et al., 2018). Adenosine triphosphate (ATP) loss may be a reason for increased muscle stiffness (Spudich, 2001), because ATP is required to detach myosin from actin during muscle contraction cy-

cles. With ATP loss during repeated exercise, this detachment ability declines, and the two proteins remain connected (Spudich, 2001). However, in this study, DS followed by VFR significantly decreased muscle stiffness, potentially reducing the risk of sports injury (e.g., muscle strain). We speculate that the following mechanism underlies the reduced muscle stiffness after VFR: rollers may modulate myofascial tone through changes in thixotropic properties, blood flow, and fascial hydration, affecting tissue stiffness (Behm and Wilke, 2019). In addition to these effects, the transmission of additional mechanical oscillations to the target muscles affects several physiological systems, such as joint mechanoreceptors (e.g., the Golgi tendon organ) (Lee et al., 2018; Moezy et al., 2008). Some limitations of the study warrant mention. First, only DS and combined DS and VFR warm-ups were conducted and compared and not static stretching or VFR alone. Different combinations may have other effects. Second, whether VFR prior to DS is more effective must be verified in the future. Third, the density of outer foam layers of rollers can differ and may affect outcomes (Cheatham and Stull, 2019).

Conclusion

The findings suggest that DS with or without additional VFR as a warm-up protocol significantly improves ROM, CMJ height, and agility. In addition, VFR considerably offset muscle stiffness compared with DS alone. Therefore, we suggest that practitioners consider DS as a first line of warm-up exercise to increase ROM, CMJ height, and agility in badminton athletes. Moreover, the addition of VFR to DS results in a large reduction of muscle stiffness, which may be advantageous for improving exercise performance and reducing the risk of injury. The findings may also be useful in athletic practice settings. Athletes, coaches and athletic professionals may consider them when selecting effective warm-up practices to augment athletic performance.

Acknowledgements

We would like to thank all our participants for their time and effort. We also thank researcher for assistance with statistical analyses, from the Division of Medical Statistics and Bioinformatics, Department of Medical Research, Kaohsiung Medical University Hospital, Kaohsiung Medical University. This study was supported by the NSYSU-KMU as joint research projects (NSYSUKMU 107-P021; NSYSUKMU 109-P005). The experiments comply with the current laws of the country in which they were performed. The authors have no conflict of interest to declare.

References

- Aird, L., Samuel, D. and Stokes, M. (2012) Quadriceps muscle tone, elasticity and stiffness in older males: reliability and symmetry using the MyotonPRO. *Archives of Gerontology and Geriatrics* **55**, e31-e39.
- Balsalobre-Fernández, C., Glaister, M. and Lockey, R.A. (2015) The validity and reliability of an iPhone app for measuring vertical jump performance. *Journal of Sports Sciences* **33**, 1574-1579.
- Behm, D.G. and Chaouachi, A. (2011) A review of the acute effects of static and dynamic stretching on performance. *European Journal of Applied Physiology* **111**, 2633-251.
- Behm, D.G. and Wilke, J. (2019) Do Self-Myofascial Release Devices Release Myofascia? Rolling Mechanisms: A Narrative Review. *Sports Medicine* **49**, 1173-1181.
- Chaouachi, A., Castagna, C., Chtara, M., Brughelli, M., Turki, O., Galy, O., Chamari, K. and Behm, D.G. (2010) Effect of warm-ups involving static or dynamic stretching on agility, sprinting, and jumping performance in trained individuals. *The Journal of Strength and Conditioning Research* **24**, 2001-2011.
- Chatzopoulos, D., Galazoulas, C., Patikas, D. and Kotzamanidis, C. (2014) Acute effects of static and dynamic stretching on balance, agility, reaction time and movement time. *Journal of Sports Science and Medicine* **13**, 403-409.
- Cheatham, S.W. and Stull, K.R. (2019) Roller massage: Comparison of three different surface type pattern foam rollers on passive knee range of motion and pain perception. *Journal of Bodywork and Movement Therapies* **23**, 555-560.
- Cheatham, S.W., Stull, K.R. and Kolber, M.J. (2019) Comparison of a vibration roller and a nonvibration roller intervention on knee range of motion and pressure pain threshold: a randomized controlled trial. *Journal of Sport Rehabilitation* **28**, 39-45.
- Chen, C.H., Xin, Y., Lee, K.W., Lin, M.J. and Lin, J.J. (2018) Acute effects of different dynamic exercises on hamstring strain risk factors. *PLoS One* **13**, e0191801.
- Chen, S.M., Shen, F.C., Chen, J.F., Chang, W.D. and Chang, N.J. (2019) Effects of Resistance Exercise on Glycated Hemoglobin and Functional Performance in Older Patients with Comorbid Diabetes Mellitus and Knee Osteoarthritis: A Randomized Trial. *International Journal of Environmental Research and Public Health* **17**, 224.
- Cochrane, D.J. (2011) The potential neural mechanisms of acute indirect vibration. *Journal of Sports Science and Medicine* **10**, 19-30.
- Costa, P.B., Herda, T.J., Herda, A.A. and Cramer, J.T. (2014) Effects of dynamic stretching on strength, muscle imbalance, and muscle activation. *Medicine & Science in Sports & Exercise* **46**, 586-593.
- De Benito, A.M., Valldecabres, R., Ceca, D., Richards, J., Igual, J.B. and Pablos, A. (2019) Effect of vibration vs non-vibration foam rolling techniques on flexibility, dynamic balance and perceived joint stability after fatigue. *PeerJ* **7**, e8000.
- Ditroilo, M., Hunter, A.M., Haslam, S. and De Vito, G. (2011) The effectiveness of two novel techniques in establishing the mechanical and contractile responses of biceps femoris. *Physiological Measurement* **32**, 1315-1326.
- Faul, F., Erdfelder, E., Lang, A.G. and Buchner, A. (2007) G*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods* **39**, 175-191.
- Fletcher, I.M. (2010) The effect of different dynamic stretch velocities on jump performance. *European Journal of Applied Physiology* **109**, 491-498.
- Garcia-Gutierrez, M.T., Guillen-Rogel, P., Cochrane, D.J. and Marin, P.J. (2018) Cross transfer acute effects of foam rolling with vibration on ankle dorsiflexion range of motion. *Journal of Musculoskeletal and Neuronal Interactions* **18**, 262-267.
- Herda, T.J., Herda, N.D., Costa, P.B., Walter-Herda, A.A., Valdez, A.M. and Cramer, J.T. (2013) The effects of dynamic stretching on the passive properties of the muscle-tendon unit. *Journal of Sports Sciences* **31**, 479-487.
- Hodgson, M., Docherty, D. and Robbins, D. (2005) Post-activation potentiation: underlying physiology and implications for motor performance. *Sports Medicine* **35**, 585-595.
- Iwata, M., Yamamoto, A., Matsuo, S., Hatano, G., Miyazaki, M., Fukaya, T., Fujiwara, M., Asai, Y. and Suzuki, S. (2019) Dynamic Stretching Has Sustained Effects on Range of Motion and Passive Stiffness of the Hamstring Muscles. *Journal of Sports Science and Medicine* **18**, 13-20.
- Kuntze, G., Mansfield, N. and Sellers, W. (2010) A biomechanical analysis of common lunge tasks in badminton. *Journal of Sports Sciences* **28**, 183-191.
- Lee, C.-L., Chu, I.-H., Lyu, B.-J., Chang, W.-D. and Chang, N.-J. (2018) Comparison of vibration rolling, nonvibration rolling, and static stretching as a warm-up exercise on flexibility, joint proprioception, muscle strength, and balance in young adults. *Journal of Sports Sciences* **36**, 2575-2582.
- Lees, A. (2003) Science and the major racket sports: a review. *Journal of Sports Sciences* **21**, 707-732.
- Lim, J.-H., Park, C.-B. and Kim, B.-G. (2019) The effects of vibration foam roller applied to hamstring on the quadriceps electromyography activity and hamstring flexibility. *Journal of Exercise Rehabilitation* **15**, 560-565.

- Lim, J.H. and Park, C.B. (2019) The immediate effects of foam roller with vibration on hamstring flexibility and jump performance in healthy adults. *Journal of Exercise Rehabilitation* **15**, 50-54.
- Lyu, B.J., Lee, C.L., Chang, W.D. and Chang, N.J. (2020) Effects of Vibration Rolling with and without Dynamic Muscle Contraction on Ankle Range of Motion, Proprioception, Muscle Strength and Agility in Young Adults: A Crossover Study. *International Journal of Environmental Research and Public Health* **17**, 354
- MacDonald, G.Z., Penney, M.D., Mullaley, M.E., Cuconato, A.L., Drake, C.D., Behm, D.G. and Button, D.C. (2013) An acute bout of self-myofascial release increases range of motion without a subsequent decrease in muscle activation or force. *The Journal of Strength & Conditioning Research* **27**, 812-821.
- McMillian, D.J., Moore, J.H., Hatler, B.S. and Taylor, D.C. (2006a) Dynamic vs. static-stretching warm up: the effect on power and agility performance. *The Journal of Strength & Conditioning Research* **20**, 492-499.
- McMillian, D.J., Moore, J.H., Hatler, B.S. and Taylor, D.C. (2006b) Dynamic vs. static-stretching warm up: the effect on power and agility performance. *The Journal of Strength & Conditioning Research* **20**, 492-429.
- Mei, Q., Gu, Y., Fu, F. and Fernandez, J. (2017) A biomechanical investigation of right-forward lunging step among badminton players. *Journal of Sports Sciences* **35**, 457-462.
- Miyamoto, N., Hirata, K. and Kanehisa, H. (2017) Effects of hamstring stretching on passive muscle stiffness vary between hip flexion and knee extension maneuvers. *Scandinavian Journal of Medicine & Science in Sports* **27**, 99-106.
- Moezy, A., Olyaci, G., Hadian, M., Razi, M. and Faghizadeh, S. (2008) A comparative study of whole body vibration training and conventional training on knee proprioception and postural stability after anterior cruciate ligament reconstruction. *British Journal of Sports Medicine* **42**, 373-378.
- Nordez, A., McNair, P., Casari, P. and Cornu, C. (2009) The effect of angular velocity and cycle on the dissipative properties of the knee during passive cyclic stretching: a matter of viscosity or solid friction. *Clinical Biomechanics* **24**, 77-81.
- Opar, D.A., Williams, M.D. and Shield, A.J. (2012) Hamstring strain injuries: factors that lead to injury and re-injury. *Sports Medicine* **42**, 209-226.
- Opplert, J. and Babault, N. (2018) Acute effects of dynamic stretching on muscle flexibility and performance: an analysis of the current literature. *Sports Medicine* **48**, 299-325.
- Pamboris, G.M., Noorkoiv, M., Baltzopoulos, V., Gokalp, H., Marzilger, R. and Mohagheghi, A.A. (2018) Effects of an acute bout of dynamic stretching on biomechanical properties of the gastrocnemius muscle determined by shear wave elastography. *PLoS One* **13**, e0196724.
- Paradisis, G.P., Pappas, P.T., Theodorou, A.S., Zacharogiannis, E.G., Skordilis, E.K. and Smirmiotou, A.S. (2014) Effects of static and dynamic stretching on sprint and jump performance in boys and girls. *The Journal of Strength & Conditioning Research* **28**, 154-160.
- Perrier, E.T., Pavol, M.J. and Hoffman, M.A. (2011) The acute effects of a warm-up including static or dynamic stretching on countermovement jump height, reaction time, and flexibility. *The Journal of Strength & Conditioning Research* **25**, 1925-1931.
- Piva, S.R., Fitzgerald, K., Irrgang, J.J., Jones, S., Hando, B.R., Browder, D.A. and Childs, J.D. (2006) Reliability of measures of impairments associated with patellofemoral pain syndrome. *BMC Musculoskeletal Disorders* **7**, 33.
- Pruyn, E.C., Watsford, M.L. and Murphy, A.J. (2016) Validity and reliability of three methods of stiffness assessment. *Journal of Sport and Health Science* **5**, 476-483.
- Romero-Moraleda, B., González-García, J., Cuéllar-Rayó, Á., Balsalobre-Fernández, C., Muñoz-García, D. and Morencos, E. (2019) Effects of vibration and non-vibration foam rolling on recovery after exercise with induced muscle damage. *Journal of Sports Science and Medicine* **18**, 172-180.
- Ryan, E.D., Everett, K.L., Smith, D.B., Pollner, C., Thompson, B.J., Sobolewski, E.J. and Fiddler, R.E. (2014a) Acute effects of different volumes of dynamic stretching on vertical jump performance, flexibility and muscular endurance. *Clinical Physiology and Functional Imaging* **34**, 485-492.
- Ryan, E.D., Everett, K.L., Smith, D.B., Pollner, C., Thompson, B.J., Sobolewski, E.J. and Fiddler, R.E. (2014b) Acute effects of different volumes of dynamic stretching on vertical jump performance, flexibility and muscular endurance. *Clinical Physiology and Functional Imaging* **34**, 485-492.
- Spudich, J.A. (2001) The myosin swinging cross-bridge model. *Nature Reviews Molecular Cell Biology* **2**, 387-392.
- Stanton, R., Kean, C.O. and Scanlan, A.T. (2015) My Jump for vertical jump assessment. *British Journal of Sports Medicine* **49**, 1157-1158.
- Su, H., Chang, N.-J., Wu, W.-L., Guo, L.-Y. and Chu, I.-H. (2017) Acute effects of foam rolling, static stretching, and dynamic stretching during warm-ups on muscular flexibility and strength in young adults. *Journal of Sport Rehabilitation* **26**, 469-477.
- Tiwari, L., Rai, V. and Srinet, S. (2011) Relationship of selected motor fitness components with the performance of badminton player. *Asian Journal of Physical Education and Computer Science in Sports* **5**, 88-91.
- Turki, O., Chaouachi, A., Drinkwater, E.J., Chtara, M., Chamari, K., Amri, M. and Behm, D.G. (2011) Ten minutes of dynamic stretching is sufficient to potentiate vertical jump performance characteristics. *The Journal of Strength & Conditioning Research* **25**, 2453-2463.
- Turki, O., Dhahbi, W., Padulo, J., Khalifa, R., Ridene, S., Alamri, K., Milic, M., Gueid, S. and Chamari, K. (2019) Warm-Up With Dynamic Stretching: Positive Effects on Match-Measured Change of Direction Performance in Young Elite Volleyball Players. *International Journal of Sports Physiology and Performance* **6**, 1-6.
- Wong, T.K.K., Ma, A.W.W., Liu, K.P.Y., Chung, L.M.Y., Bae, Y.H., Fong, S.S.M., Ganesan, B. and Wang, H.K. (2019) Balance control, agility, eye-hand coordination, and sport performance of amateur badminton players: A cross-sectional study. *Medicine (Baltimore)* **98**, e14134.
- Woods, K., Bishop, P. and Jones, E. (2007) Warm-up and stretching in the prevention of muscular injury. *Sports Medicine* **37**, 1089-1099.
- Youdas, J.W., Krause, D.A., Hollman, J.H., Harmsen, W.S. and Laskowski, E. (2005) The influence of gender and age on hamstring muscle length in healthy adults. *Journal of Orthopaedic & Sports Physical Therapy* **35**, 246-252.

Key points

- Dynamic stretching (DS) with or without vibration foam rolling (VFR) significantly increased knee range of motion, jump height, and agility
- However, DS increased thigh and calf muscle stiffness.
- DS followed by VFR significantly decreased muscle stiffness, potentially reducing the risk of sports injury.

AUTHOR BIOGRAPHY



Wei-Cheng LIN

Employment

Department of Sports Medicine at Kaohsiung Medical University

Degree

MS

Research interests

Sports medicine, Physical therapy

E-mail: diekid0910@gmail.com

	<p>Chia-Lun LEE Employment Full professor in Center for Physical and Health Education at National Sun Yat-sen University Degree PhD Research interests Strength and conditioning training, Sport nutrition, Health promotion E-mail: karenlee1129@gmail.com</p>
	<p>Nai-Jen CHANG Employment Associate professor in Department of Sports Medicine at Kaohsiung Medical University Degree PhD Research interests Sports medicine, Cartilage repair, Health promotion E-mail: njchang@kmu.edu.tw</p>

✉ **Nai-Jen Chang**

Institution: Department of Sports Medicine, Kaohsiung Medical University, 100 Shih-Chuan 1st Rd., Kaohsiung City 807, Taiwan