

Formation and Function of the Myofibroblast during Tissue Repair

Boris Hinz¹

It is generally accepted that fibroblast-to-myofibroblast differentiation represents a key event during wound healing and tissue repair. The high contractile force generated by myofibroblasts is beneficial for physiological tissue remodeling but detrimental for tissue function when it becomes excessive such as in hypertrophic scars, in virtually all fibrotic diseases and during stroma reaction to tumors. Specific molecular features as well as factors that control myofibroblast differentiation are potential targets to counteract its development, function, and survival. Such targets include α -smooth muscle actin and more recently discovered markers of the myofibroblast cytoskeleton, membrane surface proteins, and the extracellular matrix. Moreover, intervening with myofibroblast stress perception and transmission offers novel strategies to reduce tissue contracture; stress release leads to the instant loss of contraction and promotes apoptosis.

Journal of Investigative Dermatology (2007) **127**, 526–537. doi:10.1038/sj.jid.5700613

Introduction

It is generally accepted that myofibroblasts represent key players in the physiological reconstruction of connective tissue after injury and in generating the pathological tissue deformations that characterize fibrosis (Gabbiani, 2003; Desmouliere *et al.*, 2005). In previous reviews, we have focused on the role of cell-cell and cell-matrix contacts in myofibroblast development and function (Hinz and Gabbiani, 2003a, b; Hinz, 2006). Here, I draw a more general picture how the mechanical and chemical microenvironments integrate to promote tissue remodeling by myofibroblast. Differentiation of fibroblasts into myofibroblasts can be understood as a two-step process: (1) to re-populate damaged tissues, fibroblasts acquire a migratory phenotype by *de novo* developing contractile bundles. These *in vivo* stress fibers are first composed of cytoplasmic actins and generate comparably small traction forces (Hinz *et al.*, 2001b). We have recently proposed the term “proto-myofibroblast” to discriminate such

activated fibroblasts from quiescent fibroblasts that are devoid of any contractile apparatus in most intact tissues (Tomasek *et al.*, 2002). This first phenotypic change occurs in response to changes in the composition, organization, and mechanical property of the extracellular matrix (ECM) (Hinz and Gabbiani, 2003b) and to cytokines locally released by inflammatory and resident cells (Werner and Grose, 2003). (2) With increasing stress in the ECM resulting from their own remodeling activity, proto-myofibroblasts further develop into “differentiated myofibroblasts” by neo-expressing α -smooth muscle actin (α -SMA), the most widely used myofibroblast marker. Expression of α -SMA is precisely controlled by the joint action of growth factors like transforming growth factor (TGF β 1), of specialized ECM proteins like the fibronectin (FN) splice variant ED-A FN, and of the mechanical microenvironment (Tomasek *et al.*, 2002). Incorporation of α -SMA into stress fibers significantly augments the contractile activity of fibroblastic cells

and hallmarks the contraction phase of connective tissue remodeling (Hinz *et al.*, 2001a).

It has to be noted that the contribution of myofibroblast contraction to physiological tissue remodeling has been questioned by a study reporting normal closure of pig full thickness wounds despite repeated excision of the central and peripheral granulation tissue (Gross *et al.*, 1995). However, rat wounds that were kept open for 10 days with a plastic frame and have then been released from the splint contract to ~50% of their initial size within 6 hours; this cannot be explained by enhanced proliferation of fibroblasts and keratinocytes at the wound edge. Hence, the porcine skin may be able to compensate for the experimentally induced chronic loss of granulation tissue by mechanisms that have not yet been elucidated. Such a compensatory mechanism appears to function in α -SMA-knockout mice that macroscopically exhibit almost normal wound closure (personal communication, J.J. Tomasek, University of Oklahoma Health

¹Laboratory of Cell Biophysics, École Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland

Correspondence: Dr Boris Hinz, Laboratory of Cell Biophysics, École Polytechnique Fédérale de Lausanne (EPFL), Bâtiment SG-AA-B 143, Station 15, CH-1015 Lausanne, Switzerland. E-mail: boris.hinz@epfl.ch

Abbreviations: α -SMA, α -smooth muscle actin; ECM, extracellular matrix; FA, focal adhesion; Fizz, found in inflammatory zone; FN, fibronectin; LAP, latency-associated protein; LTBP-1, latent TGF β 1-binding protein; SMC, smooth muscle cell; TGF, transforming growth factor

Received 29 May 2006; revised 6 July 2006; accepted 17 July 2006

Sciences Center, Oklahoma City, OK). These mice appear to substitute for α -SMA by *de novo* expressing other muscle actin isoforms like γ -smooth muscle and α -skeletal actin, which also explains why blood pressure is surprisingly little affected by the lack of α -SMA in vascular smooth muscle cells (SMCs) (Schildmeyer *et al.*, 2000). In contrast to these chronic conditions, acute inhibition of myofibroblast contraction by intracellular delivery of the specific N-terminal sequence AcEEED of α -SMA significantly reduces wound contraction (Hinz *et al.*, 2002). It will be interesting to test whether conditional α -SMA-knockout mice or mice treated with α -SMA RNAi will exhibit similarly impaired wound closure.

In physiological remodeling such as during dermal wound healing, the contractile activity of myofibroblasts is terminated when the tissue is repaired; α -SMA expression then decreases and myofibroblasts disappear by apoptosis (Desmouliere *et al.*, 1995). In pathological wound healing, however, myofibroblast activity persists and leads to tissue deformation, which is particularly evident in hypertrophic scars developing after burn injury, in the fibrotic phase of scleroderma and in the palmar fibromatosis of Dupuytren's disease (Schurch *et al.*, 2006). Myofibroblast-generated contractures are also characteristic for fibrosis affecting vital organs such as the liver (Desmouliere *et al.*, 2003), heart (Virag and Murry, 2003; Brown *et al.*, 2005), lung (Phan, 2002; Thannickal *et al.*, 2004), and kidney (Lan, 2003). In addition, myofibroblast participation to the process called stroma reaction promotes cancer progression by creating a stimulating microenvironment for epithelial tumor cells (De Wever and Mareel, 2003; Desmouliere *et al.*, 2004). It becomes increasingly accepted that stroma cells represent important targets of anticancer treatments (Bissell and Radisky, 2001; Liotta and Kohn, 2001; Mueller and Fusenig, 2004). In the light of such severe consequences of myofibroblast appearance and dysfunction, the necessity of more profoundly understanding the molecular mechanisms of myofibro-

blast formation and function appears clear.

Myofibroblast origin

The classical view on dermal wound healing implies local recruitment of fibroblasts from the dermis of the intact adjacent skin to sites of inflammation (Desmouliere *et al.*, 2005). Pericytes and SMC from the vasculature have been proposed as another local myofibroblast source in scleroderma (Rajkumar *et al.*, 2005) as well as liver and glomerular fibrosis (Desmouliere *et al.*, 2003). In addition, fibroblasts may originate from fibrocytes, a sub-population of bone marrow-derived leukocytes with fibroblast characteristics (Abe *et al.*, 2001). This concept has recently been revisited by transplanting bone marrow from (1) male to irradiated female mice and identifying the Y chromosome by *in situ* hybridization (Direkze *et al.*, 2003) and (2) from transgenic green fluorescent mice to irradiated wild-type animals (Hashimoto *et al.*, 2004). These studies reveal a surprisingly high fraction of 30–50% of the wound myofibroblasts potentially deriving from fibrocyte progenitors (Direkze *et al.*, 2003; Ishii *et al.*, 2005; Mori *et al.*, 2005). Comparable proportions have been demonstrated for myofibroblasts appearing during fibrosis of the liver (Forbes *et al.*, 2004), kidney (Direkze *et al.*, 2003), and lung (Hashimoto *et al.*, 2004) and during the stroma reaction to epithelial tumors (Ishii *et al.*, 2003). It remains to be seen whether myofibroblasts from different origins exhibit different characteristics and functions during tissue repair as suggested for liver fibrosis (Guyot *et al.*, 2006) or whether the organism is recruiting myofibroblast precursors from several sources to satisfy the temporarily high demand of contractile cells, which may all follow a similar differentiation program.

Identification of the myofibroblast – a warrant

The question of the myofibroblast origin is closely related to the problem of its identification. Three major ultrastructural features discriminate myofibroblasts from quiescent fibroblasts in tissues: (1) bundles of contractile mi-

crofilaments, (2) extensive cell-to-matrix attachment sites, and (3) intercellular adherens and gap junctions (Eyden, 2005; Schurch *et al.*, 2006). However, this definition has its limits when myofibroblasts need to be discriminated from other contractile cell types like SMC, requiring specific molecular markers.

Cytoskeletal markers of the myofibroblast.

The most frequently employed myofibroblast marker is α -SMA, which evidently fails to distinguish between myofibroblasts and SMC in situations that exhibit mixed populations. For example, remodeling of injured arteries is thought to be predominantly driven by SMC from the media but a contribution from adventitial fibroblasts has also been suggested (Sartore *et al.*, 2001; Zalewski *et al.*, 2002). Contractile SMC specifically express smooth muscle myosin heavy chain, h-caldesmon, and desmin; however, SMC lose these markers when acquiring a synthetic phenotype and after being placed in culture (Christen *et al.*, 2001). Until recently, smoothelin was suggested a late differentiation marker for SMC that is not expressed in myofibroblasts (van der Loop *et al.*, 1996). However, gene expression profiling supported by protein biochemistry revealed induction of smoothelin and other late SMC markers in TGF β 1-treated cultured lung fibroblasts (Chambers *et al.*, 2003). Very recently, the 4Ig isoform of the stress fiber protein palladin has been proposed as novel marker for myofibroblast differentiation (Rönty *et al.*, 2006), but Western blotting analysis with pan-palladin antibodies indicates expression of this isoform also in SMC (Mykkanen *et al.*, 2001). Hence, at present no cytoskeletal protein allows to reliably discriminate between myofibroblasts and SMC; however, recent advances in proteomics and gene array analysis may lead to the identification of such a unique marker of the myofibroblast, provided that it exists (Chambers *et al.*, 2003; Malmstrom *et al.*, 2004).

The myofibroblast surface. Of particular interest not only for the purpose of

identification but also for the delivery of specific drugs are proteins specifically expressed on the plasma membrane of myofibroblasts. The membrane protein Thy-1 is frequently used for the inverse approach as it is expressed by fibroblasts but absent from differentiated myofibroblasts (Koumas *et al.*, 2003). Interestingly, Thy-1-negative cells fail to respond to fibrogenic stimuli suggesting that Thy-1 expression characterizes a specific subpopulation of fibroblasts that may not be able to undergo myofibroblast differentiation (Zhou *et al.*, 2004; Hagood *et al.*, 2005). A promising new strategy to target and identify myofibroblasts is the expression of specific cadherins (Hinz and Gabbiani, 2003a), transmembrane cell-cell adhesion proteins that are intracellularly linked to the actin cytoskeleton (Gumbiner, 2005). In contrast to virtually cadherin-negative fibroblasts of normal dermis, proto-myofibroblasts of early granulation tissue and in culture express N-cadherin (cadherin-2, A-cell adhesion molecule (A-CAM)) (Hinz *et al.*, 2004). In conjunction with the appearance of α -SMA after TGF β 1-treatment in culture or in differentiated myofibroblasts of contractile wound granulation tissue, N-cadherin becomes gradually replaced by OB-cadherin (cadherin-11). OB- but not N-cadherin seems to play a functional role in coordinating myofibroblast contraction in populations (Hinz *et al.*, 2004). A similar shift from N- to OB-cadherin expression has been described in stroma fibroblasts reacting against human prostate cancer progression (Tomita *et al.*, 2000). OB-cadherin is absent from SMC but it is expressed on a variety of cell types of mesenchymal origin; hence, it cannot be considered as unique myofibroblast surface marker.

The ECM of myofibroblast-populated tissue. In addition to producing tension for remodeling, enhanced ECM synthesizing and processing activity significantly contributes to tissue remodeling. Myofibroblasts produce several components of the ECM that can potentially be used as molecular markers, depending on the context of myofibroblast appearance; the most prominent myo-

fibroblast ECM products are collagens of types I, III, IV, and V (Schurch *et al.*, 2006) which, however, are produced by a variety of cells. More recently, collagen type VI attracted attention as it is upregulated together with myofibroblast differentiation in human renal fibrosis (Groma, 1998), in myocardial interstitial fibrosis (Kitamura *et al.*, 2001), and in the stroma of hepatocellular carcinoma (Faouzi *et al.*, 1999). Interestingly, culturing cardiac fibroblasts on collagen type VI substrates induces myofibroblast differentiation but not culture on collagen types I and III (Naugle *et al.*, 2006). At present, the most reliable marker of the myofibroblast ECM is the FN splice variant ED-A FN (Serini *et al.*, 1998). ED-A FN is also expressed in low amounts by fibroblastic cells in culture (Dugina *et al.*, 2001; Hinz *et al.*, 2001a) and by vascular SMC *in vivo* and *in vitro* (Glukhova *et al.*, 1989). Another component of the myofibroblast ECM, glycoprotein tenascin-C is associated with tissue repair phenomena (Chiquet-Ehrismann and Chiquet, 2003). Tenascin-C appears to attract fibroblasts and to promote their differentiation into myofibroblast in injured tissue (Tamaoki *et al.*, 2005) and at the tumor invasion front of cancers (De Wever *et al.*, 2004). However, the shared expression in SMC limits its usage as myofibroblast marker (Chiquet-Ehrismann and Chiquet, 2003).

In view of the expression profile of cytoskeletal and ECM proteins, the differentiated myofibroblast appears to be situated in a continuous spectrum existing between fibroblasts and SMC. It may be necessary to combine common markers of the contractile cell phenotype with the exclusive surface expression of proteins like OB-cadherin and Thy-1 to unmistakably discriminate between fibroblasts, myofibroblasts, and SMC.

Chemical factors inducing myofibroblast formation

Apart from being important diagnostic tools, myofibroblast-specific proteins are likewise potential targets to modulate myofibroblast differentiation and/or contractile function; this has been shown by interfering with α -SMA (Hinz

et al., 2002) and with ED-A FN (Serini *et al.*, 1998). Alternative antifibrosis strategies aim to prevent development of tissue contractures by affecting physiological cytokines that control myofibroblast differentiation. TGF β 1 is considered the major growth factor directly promoting myofibroblast development by inducing expression of α -SMA (Desmouliere *et al.*, 1993; Ronnov-Jessen and Petersen, 1993), ECM proteins (Werner and Grose, 2003), and a number of cytoskeletal proteins that construct the myofibroblast contractile apparatus (Malmstrom *et al.*, 2004). TGF β 1 effects are "fine-tuned" by cooperative or antagonistic growth factors and some recent studies report myofibroblast differentiation in the absence of TGF β 1 signaling.

Control of α -SMA transcription by TGF β 1 signaling.

The major pathway through which TGF β 1 regulates expression of α -SMA in fibroblastic cells appears to involve Smad signaling (Figure 1). Binding of active TGF β 1 to the TGF β receptor type II leads to the phosphorylation and recruitment of TGF β receptor type I into a heteromeric receptor complex. The serine/threonine kinase activity of the activated complex phosphorylates Smad2 and Smad3, that each bind to Smad4 and translocate into the nucleus to enhance gene transcription by cooperating with DNA transcription factors (Feng and Derynck, 2005; Massague *et al.*, 2005). Transcription of α -SMA in lung myofibroblasts and in myofibroblast-like activated hepatic stellate cells is predominantly mediated by the binding of Smad3 to the Smad-binding element 1 upstream of the α -SMA core promoter sequence; the role of Smad3 binding to a second Smad-binding element (Smad-binding element 2) in the core region is unclear at present (Hu *et al.*, 2003; Uemura *et al.*, 2005). In contrast, another study demonstrated that overexpression of Smad2 but not of Smad3 induces myofibroblast differentiation in lung fibroblasts (Evans *et al.*, 2003); this discrepancy may be due to different roles of Smad2 and Smad3 depending on the level of myofibroblast differentiation (Liu *et al.*, 2003). Moreover, rather than being essential factors for α -




Figure 1. Regulation of α -SMA transcription in myofibroblasts. The major pathway of TGF β 1-induced α -SMA expression in myofibroblasts is mediated via Smad3 activation by the TGF β 1 receptor complex, leading to Smad3 association with Smad4 and translocation into the nucleus. Smad3 binding to Smad-binding elements in the promoter region regulates α -SMA transcription in conjunction with a variety of transcription factors (TF). Some TGF β 1-antagonizing factors like IFN- γ operate via activation of YB-1 that inhibits Smad3-enhanced gene transcription and upregulates expression of the inhibitory Smad7. TGF β 1-induced, Smad-independent transcription of α -SMA is enhanced by binding of the krüppel-like factors Sp1/3 to the TGF β 1 hypersensitivity region. Moreover, binding of the TEF-1 to the MCAT-1 element is crucial for α -SMA expression in myofibroblasts and SMCs. In contrast to SMC, interaction of the serum-response factor with CARG elements A, B, C, and D is not mandatory for α -SMA transcription; however, the intronic CARG element regulates α -SMA transcription in both cell types. Please refer to text for details and references.

SMA transcription, Smad 2/3 should be considered as modulators of the activity of other transcription factors. This may explain different results obtained from acute removal of Smad, for example, by using an antisense approach compared with chronic Smad deficiency in knockout animals. In the first setting, cells may not be able to compensate for the instant loss of the important modulating activity of Smads; adaptation and compensation may, however, occur in the knockout system.

Alternatively, TGF β 1 can regulate fibroblast gene expression independently from Smad signaling (Derynck and Zhang, 2003; Moustakas and Heldin, 2005) and expression of α -SMA has been documented during the

activation of Smad3-null hepatic stellate cells in culture (Schnabl *et al.*, 2001). One pathway of Smad-independent TGF β 1 induction of the myofibroblastic phenotype has been revealed in Smad3-knockout fibroblasts, transfected with dominant-negative Smad2 and appears to function via activating phosphatidylinositol 3-kinase and the downstream effector p21-activated kinase-2; expression of α -SMA has not been assessed in this study (Wilkes *et al.*, 2005). Smad-independent transcription of α -SMA in fibroblasts is regulated via the TGF β 1 control element (Roy *et al.*, 2001; Tomasek *et al.*, 2005) that together with the TGF β 1 hypersensitivity region serves as binding site for the transcriptional activator proteins Sp1/2, members of

the family of krüppel-like factors (Cogan *et al.*, 2002; Subramanian *et al.*, 2004). Another mandatory element for α -SMA transcription in myofibroblasts and SMC is the MCAT-1 site that is enhanced by binding of the transcription enhancer factor-1 which apparently competes with MCAT-suppressor proteins; the presence of a second MCAT element further upstream of the promoter appears not crucial (Swartz *et al.*, 1998; Carlini *et al.*, 2002).

Importantly, transcription of α -SMA in SMC differs from fibroblasts by being additionally regulated via serum-response factor binding to CC(A/T)6GG-like sequence motifs (CARG elements) in the promoter region (Hautmann *et al.*, 1997); these elements seem not

to be required for α -SMA transcription in myofibroblasts (Roy *et al.*, 2001). In contrast, the CARG element in the first intron promoter region has recently been demonstrated to regulate α -SMA transcription in myofibroblasts similar to SMC (Tomasek *et al.*, 2005). It remains to be shown whether upregulated serum-response factor in response to mechanical stimulation (Wang *et al.*, 2006) acts predominantly through this intronic CARG element to enhance α -SMA expression. Deciphering the distinct upstream events in each cell type may provide future gene targeting strategies to specifically inhibit myofibroblast formation without affecting SMC function.

Soluble factors cooperating with TGF β 1 in myofibroblast differentiation. One of the best studied cytokines induced by TGF β 1 is connective tissue growth factor that potentiates the profibrotic action of TGF β 1 and at the same time induces fibroblast proliferation (Leask *et al.*, 2002). These pleiotropic effects have been attributed to distinct connective tissue growth factor subdomains and to the mutually exclusive cooperation with other cytokines like EGF and IGF-2 (Grotendorst and Duncan, 2005). Connective tissue growth factor cooperates with TGF β 1 but does not substitute for its action, as α -SMA expression remains unchanged after administration of recombinant connective tissue growth factor to cultures of corneal fibroblasts (Folger *et al.*, 2001). A novel cooperative factor of TGF β 1 seems to be the cell-surface protein galectin-3. Galectin-3 is mandatory for TGF β 1-induced myofibroblast differentiation of hepatic stellate cells in culture and is upregulated in experimental liver, kidney, and lung fibrosis (Henderson *et al.*, 2006). Knockout of galectin-3 significantly reduces hepatic fibrosis at unchanged levels of TGF β 1 and inhibits myofibroblast activation of cultured stellate cells in the presence of TGF β 1; addition of recombinant galectin-3 rescues myofibroblast differentiation. Binding of galectin-3 to growth factor receptors has been demonstrated but its cooperation with TGF β 1 is not yet understood (Henderson *et al.*, 2006).

Soluble factors that antagonize TGF β 1-induced myofibroblast differentiation. Attractive with respect to therapeutic treatment of fibrosis are growth factors that antagonize TGF β 1 and thereby reduce myofibroblast differentiation. The inflammatory mediator IL-1 efficiently inhibits TGF β 1-induced α -SMA expression in cultured dermal fibroblasts when added as recombinant protein (Shephard *et al.*, 2004b); overexpression of the intracellular IL-1 receptor antagonist (IL-1ra) has similar effects (Kanangat *et al.*, 2006). Co-culture experiments with fibroblasts suggest that time-controlled secretion of IL-1 by keratinocytes may be one mechanism during cutaneous wound healing to suppressing fibroblast-to-myofibroblast transition in early wounds (Shephard *et al.*, 2004b). IFN- γ , a cytokine produced by T cells also reduces α -SMA expression in cultured fibroblasts. This effect appears to be mediated by activation of the repressor protein YB-1, which translocates to the nucleus and interferes with Smad3-mediated transcription of TGF β 1-induced genes (Higashi *et al.*, 2003) (Figure 1). In addition, YB-1 enhances the transcription of Smad-7, a negative regulator of Smad3 signaling (Dooley *et al.*, 2006).

Myofibroblast differentiation without TGF β 1? Only few factors have been identified that potentially induce myofibroblast differentiation independently from TGF β 1. IL-6 has been suggested to be such a factor as IL-6-null mice exhibit impaired cutaneous wound healing and reduced myofibroblast differentiation at unchanged levels of TGF β 1. Moreover, recombinant IL-6 increases α -SMA transcription in fibroblast cultures from IL-6-null mice in the absence of exogenous TGF β 1 (Gallucci *et al.*, 2006). Similarly, nerve growth factor enhances α -SMA expression in cultured skin and lung fibroblasts (Micerca *et al.*, 2001). However, as myofibroblast differentiation in response to IL-6 and nerve growth factor was not tested with simultaneous inhibition of TGF β 1 signaling, cooperation with autocrine TGF β 1 cannot be ruled out. Very recently, Fizz1 (found in inflammatory zone), a novel protein secreted

by alveolar epithelial cells, has been demonstrated to undergo dramatic upregulation in an animal model of lung fibrosis. When added to cultured fibroblasts, recombinant Fizz1 induces expression of α -SMA in the absence of TGF β 1; the molecular mechanism of Fizz1 action remains elusive (Liu *et al.*, 2004). In addition to proinflammatory cytokines, agonists of myofibroblast contraction, like angiotensin-II (Swaney *et al.*, 2005), endothelin-1 (Shi-Wen *et al.*, 2004), and thrombin (Bogatkevich *et al.*, 2003) induce expression of α -SMA expression *in vitro*; however, it is unclear whether endogenous TGF β 1 contributes to this effect. It is also conceivable that contraction agonists enhance α -SMA expression by augmenting the level of intracellular stress (Shephard *et al.*, 2004a).

Myofibroblasts are controlled by the mechanical microenvironment

Despite high levels of active TGF β 1 in early wounds (Yang *et al.*, 1999), expression of α -SMA and enhanced myofibroblast contraction are restricted to the later phases of dermal wound healing. Earlier myofibroblast appearance may be suppressed by antagonistic factors released by inflammatory cells and keratinocytes. Moreover, it becomes increasingly clear that myofibroblast development is profoundly influenced by the mechanical microenvironment, in particular, by the organization and stiffness of the ECM (Hinz and Gabbiani, 2003a). The level of α -SMA expression in cultured differentiated myofibroblasts increases with increasing matrix rigidity as demonstrated using collagen and polyacrylamide gel substrates of varying stiffness (Arora *et al.*, 1999; Hinz *et al.*, 2003). Mechanically preventing wound closure by splinting the edges of experimental wounds accelerates expression of α -SMA compared with normally healing wounds; stress release by removing the splint leads to reduced α -SMA expression (Hinz *et al.*, 2001b). TGF β 1 is always present in these conditions; however, the mechanical characteristics of the environment appear to modulate its action to induce either migration or contraction (Grin-

nell and Ho, 2002). In addition to wound healing and fibrosis, altered matrix stiffness is increasingly considered as major factor activating stroma myofibroblasts in the vicinity of "stiff" tumors and thus feeding back on tumor progression (Paszek *et al.*, 2005).

What mechanical properties display the ECM in a physiological tissue context and how do they change during pathological contracture? During embryonic development, soft tissues are established with an elastic modulus of 100–20,000 Pa, depending on their location and function in the adult organism (Fung, 1993; Discher *et al.*, 2005); fibroblasts residing in intact tissues are generally "stress-shielded" by this ECM (Tomasek *et al.*, 2002). When the protective matrix structure is lost by injury, fibroblasts experience a dramatically changed mechanical microenvironment and endeavor to re-establish ECM tension by increasing remodeling and synthesizing activity (Eckes and Krieg, 2004; Marenzana *et al.*, 2006). The stiffness of the provisional ECM of early wounds is comparable with the elastic modulus of ~10–100 Pa of newly polymerized collagen gels (Kaufman *et al.*, 2005), frequently used *in vitro* models of tissue repair (Grinnell, 2003; Carlson and Longaker, 2004). Fibroblasts initiated in such gels are devoid of contractile fibers and organize actin filaments in the cortex of the cell body and of dendritic extensions (Tamariz and Grinnell, 2002). In mechanically restrained gels, tension is gradually increasing and first induces the formation of α -SMA-negative stress fibers (Tamariz and Grinnell, 2002; Marenzana *et al.*, 2006); this is similar to the *de novo* appearance of proto-myofibroblast in 5- to 6-day-old rat wound granulation tissue (Hinz *et al.*, 2001b). Gel and tissue stiffness at this stage have not been measured but neof ormation of stress fibers in fibroblasts grown on soft two-dimensional culture substrates occurs at an elastic modulus of ~3,000–6,000 Pa (Discher *et al.*, 2005; Yeung *et al.*, 2005).

Progression to the differentiated myofibroblast requires significantly higher stiffness that begins to develop in anchored collagen gels after 2–3

days of remodeling by proto-myofibroblasts (Hinz, 2006) and after 8–9 days in experimental rat wounds (Hinz *et al.*, 2001b). The threshold stiffness for *de novo* expression of α -SMA in stress fibers ranges around 20,000 Pa as demonstrated for contractile wound granulation tissue and for myofibroblasts cultured on elastic substrates (Goffin *et al.*, 2006). A comparable matrix stiffness of ~15,000 Pa activates hepatic stellate cells into α -SMA-positive myofibroblasts *in vitro* and during development of liver fibrosis *in vivo* (Wells, 2005). In other fibrotic tissues and in granulation tissue toward the end of wound healing, a matrix stiffness of greater than 50,000 Pa has been measured (Goffin *et al.*, 2006); this significant tissue rigidification by fibrogenic stroma cells is used to localize tumor metastases *in situ* with improving imaging methods (Greenleaf *et al.*, 2003).

The mechanisms and intracellular pathways through which tension potentially controls α -SMA transcription have recently been reviewed (Wang *et al.*, 2006). In addition, α -SMA itself is beginning to be considered as mechano-sensitive protein that only localizes to stress fibers under significant mechanical load; this provides a rapid mechanism to control myofibroblast contractile function (Goffin *et al.*, 2006; Hinz, 2006). Reducing stress fiber tension by reducing substrate stiffness and by inhibiting myosin contraction leads to the disassembly of α -SMA from stress fibers that persist as β -cytoplasmic actin filament bundles. Stress fiber-derived α -SMA accumulates in cytosolic aggregates that resemble short rods (Goffin *et al.*, 2006). Similar rod-like aggregates of α -SMA have been demonstrated during physiological spreading of suspended myofibroblasts (Clement *et al.*, 2005), which is characterized by gradually increasing intracellular tension and *de novo* formation of stress fibers (Hinz *et al.*, 2003). From these studies, it has been suggested that construction of α -SMA-positive stress fibers requires pre-formation of β -cytoplasmic actin bundles; absence of such an organization template seems to lead to cytosolic accumulation of α -SMA. It is conceiva-

ble that the α -SMA-specific N-terminal sequence AcEEED contributes to α -SMA mechano-sensitivity as cytoplasmic delivery of this sequence as a peptide similarly induces α -SMA rod formation (Clement *et al.*, 2005) and selectively removes α -SMA from stress fibers (Chaponnier *et al.*, 1995; Hinz *et al.*, 2002).

How myofibroblasts feel stress

The question remains how myofibroblasts sense the gradually increasing change in matrix stiffness that accompanies tissue remodeling. Three major mechanisms of mechano-sensing are conceivable: (1) mechano-sensitive ion channels in the plasma membrane (Martinac, 2004), (2) integrin-mediated stress perception (Katsumi *et al.*, 2004), and (3) geometry changes that reveal cryptic signaling domains in proteins of the ECM (Vogel and Sheetz, 2006). Cultured fibroblasts modulate the level of α -SMA expression after locally stretching the plasma membrane with the use of twisting magnetite beads; reaction to these rapid stimuli is thought to depend at least in part on the opening of mechano-sensitive ion channels and influx of Ca^{2+} over the plasma membrane (Ko *et al.*, 2001). However, myofibroblasts *in vivo* differentiate in response to matrix changes that occur sometimes over days and months, which demands a "slow" mechanosensor. Moreover, fibroblast-to-myofibroblast transition can be driven by subtle forces as produced by very slow interstitial flow that accompanies inflammation and tissue regeneration (Ng *et al.*, 2005); these physiological mechano-stimuli likely exclude membrane channels as force detection mechanism for myofibroblast differentiation.

The role of cell-matrix adhesions in myofibroblast stress perception.

It is widely accepted that to perceive mechanical signals from the ECM, fibroblasts utilize integrin-based matrix adhesions, that is, the same transmembrane organelles that anchor stress fibers to the substrate (Chen *et al.*, 2004; Bershadsky *et al.*, 2006). Ultrastructural analysis of myofibroblasts in fibrotic and wound granulation tissue

has revealed extensive cell-matrix contacts, called fibronexus that are absent from normal connective tissue fibroblasts (Eyden, 2005). Analogously, cultured myofibroblasts develop specialized focal adhesions (FAs) that have been termed "supermature FAs" to account for their significantly longer appearance (8–30 μm) (Dugina *et al.*, 2001) compared with "classical FAs" (2–6 μm) of α -SMA-negative fibroblasts (Geiger *et al.*, 2001). In addition, supermature FAs exhibit a specific molecular composition by co-expressing high levels of vinculin, paxillin, and tensin and integrins $\alpha\text{v}\beta\text{3}$ and $\alpha\text{5}\beta\text{1}$ (Dugina *et al.*, 2001; Hinz *et al.*, 2003). This is different from classical FAs that generally lack tensin and $\alpha\text{5}\beta\text{1}$ integrin and to fibrillar adhesions that contain tensin and $\alpha\text{5}\beta\text{1}$ integrin but not vinculin, paxillin, and $\alpha\text{v}\beta\text{3}$ integrin (Geiger *et al.*, 2001). More comprehensive reviews of the molecular components of myofibroblast ECM adhesions have recently been published (Hinz and Gabbiani, 2003a; Hinz, 2006). Moreover, supermature FAs display highly phosphorylated focal adhesion kinase and paxillin, which may play a role in the recruitment of fibrillar adhesion components to classical FAs (Goffin *et al.*, 2006). Consistently, phosphorylated focal adhesion kinase is central in the adhesion-dependent differentiation of cultured lung (Thannickal *et al.*, 2003) and scleroderma myofibroblast (Mimura *et al.*, 2005). It remains to be shown whether and how supermature FAs assemble in focal adhesion kinase-null corneal fibroblasts that apparently acquire a constitutive myofibroblast phenotype (Greenberg *et al.*, 2006).

Establishment of supermature FAs depends both on the significant intracellular contractile activity generated by α -SMA and on a stress-resistant ECM (Hinz, 2006). Differentiated myofibroblasts convert supermature FAs into small classical FAs after administration of the AcEEED peptide (Hinz *et al.*, 2003) and when cultured on soft substrates and in newly polymerized collagen gels (Goffin *et al.*, 2006; Hinz, 2006). These recent studies suggest that fibroblasts gain information about the mechanical state of the ECM by assessing the level of tension in stress fibers,




Figure 2. FA size controls recruitment of α -SMA to stress fibers. Arrays of ECM islets with a size of (a) $1 \times 1.25 \mu\text{m}$, (b) $4 \times 1.25 \mu\text{m}$, and (c) $10 \times 1.25 \mu\text{m}$ were created on glass by microcontact printing; non-printed regions were passivated against cell attachment (Goffin *et al.*, 2006). After 12 hours, culture differentiated myofibroblasts were stained for vinculin (red), β -cytoplasmic actin (green), and α -SMA (blue). Bar = 20 μm . Stress fibers are formed on all substrates; however, α -SMA is only recruited to contractile fibers on islets that exceed the length of classical FAs ($\leq 6 \mu\text{m}$) and exhibit supermature FA features ($> 8 \mu\text{m}$).

which is limited by the size of their matrix anchors. Forcing myofibroblasts to grow on arrays of classical FA adhesion islets ($\leq 6 \mu\text{m}$) leads to the rapid loss of α -SMA (Figure 2); dynamically enlarging these islets on extendable membranes leads to reincorporation of α -SMA into stress fibers. Reincorporation is independent from the magnitude of the applied stretch but requires that the resulting islet size corresponds to supermature FAs ($\sim 8 \mu\text{m}$). Resizing classical to supermature FAs permits exertion of ~ 4 -fold greater stress (Goffin *et al.*, 2006) and provides an exceptionally strong ECM anchor that renders myofibroblasts highly adhesive (Hinz *et al.*, 2003).

All these data suggest that formation of supermature FAs represents a critical control point in the mechanical feedback loop of extracellular stress and intracellular tension that regulates the state of myofibroblast differentiation

(Figure 3). In physiological tissue repair, this checkpoint could ensure that enhanced myofibroblast contraction only occurs when the tissue has been sufficiently remodeled by proto-myofibroblasts for effective force exertion by differentiated myofibroblasts. It appears difficult to establish whether the primary event is represented by α -SMA-generated stress fiber tension or FA supermaturation. Nevertheless, specifically targeting supermature FAs is one potential strategy to interrupt the mechanical cycle when it has become vicious in pathological contractures.

Stress-dependent conformation of ECM proteins: where mechanical and growth factor signals may converge.

The mechanical state of the ECM may directly influence myofibroblast differentiation by revealing cryptic epitopes in ECM proteins under stress. A potential candidate for such a mechanism is cellular FN, which contains cryptic




Figure 3. The mechanical feedback loop in myofibroblast development. Fibroblasts in intact tissue are stress-shielded by a functional ECM; they do not develop contractile features and cell-matrix adhesions. After injury, inflammatory signals activate fibroblasts to spread into the provisional wound matrix. Local cell remodeling activity leads to gradual increase in global matrix stiffness that counteracts cell traction forces. The resulting formation of small FAs and stress fibers that contain only cytoplasmic actins characterize the proto-myofibroblast. TGF β 1 stimulates proto-myofibroblasts to express α -SMA, which at first is not incorporated into stress fibers but organizes in cytoplasmic rod-like structures. Continuing ECM fiber alignment creates larger surfaces for adhesion formation; larger adhesions permit development of stronger stress fibers and generation of higher contractile forces. When adhesion sites grow to the size of supermature FAs, intracellular tension reaches a critical level that allows incorporation of α -SMA into pre-existing stress fibers. The force generated by α -SMA-containing stress fiber is significantly higher compared to cytoplasmic actin stress fibers leading to further FA supermaturation and ECM contraction, thereby establishing a mechanical loop. Myofibroblasts may exit this cycle when the original structure of the ECM is reconstituted and again takes over the mechanical load; stress-released myofibroblasts eventually undergo apoptosis.

domains in several type III modules that unfold upon cell traction (Vogel and Sheetz, 2006). ED-A, one of such modules is spliced into the FN molecule during tissue repair and is a prerequisite for TGF β 1-induced myofibroblast differentiation (Serini *et al.*, 1998). It remains to be shown whether cell-generated mechanical forces may change the availability of this domain for specific integrins to induce myofibroblast-promoting intracellular signals.

Another intriguing hypothesis on how increased ECM tension and/or cell traction forces enhance expression of α -SMA is by directly activating TGF β 1.

Fibroblasts and myofibroblasts secrete TGF β 1 as part of a large latent complex, additionally consisting of the latency-associated protein (LAP) and the latent TGF β 1-binding protein (LTBP-1) (Mangasser-Stephan *et al.*, 2001; Koli *et al.*, 2005). LTBP-1, a member of the fibrillin family strongly links the large latent complex to the ECM via several ECM binding sites (Unsold *et al.*, 2001). Activation of TGF β 1 occurs upon its release from the large latent complex through proteolytic cleavage (e.g. by plasmin) and by the action of thrombospondin-1 (Annes *et al.*, 2003). In epithelial cells, cell-generated traction via integrins has

been proposed as alternative mechanism to activate TGF β 1 from the large latent complex. Both, LAP and LTBP-1 contain RGD motifs for the binding of integrins (Keski-Oja *et al.*, 2004) and the epithelial integrin α v β 6 was shown to activate TGF β 1 during development of lung fibrosis and in culture independently of protease activity (Munger *et al.*, 1999; Annes *et al.*, 2004). Fibroblasts do not express α v β 6 integrin; however, integrins α v β 5 and α v β 3 were shown to bind to LAP-TGF β 1 (Ludbrook *et al.*, 2003) and both are involved in TGF β 1 activation by fibrogenic systemic sclerosis fibroblasts through a yet unknown mechanism (Asano *et al.*, 2005a,b). Moreover, the integrin α 8 β 1 was recently shown to strongly interact with LAP-TGF β 1 (Lu *et al.*, 2002) and it is upregulated in conjunction with myofibroblast differentiation during heart (Bouzeghrane *et al.*, 2004), pulmonary, and hepatic fibrosis (Levine *et al.*, 2000).

TGF β 1 activation by α v β 6 integrin requires ECM binding of LTBP-1 through its N-terminal hinge region (Annes *et al.*, 2004), suggesting that matrix resistance needs to counteract cell traction in the activation process. Consistently, transformed fibroblasts expressing high levels of latent LAP-TGF β 1 do not promote fibrosis when implanted into mouse dermis (Campagner *et al.*, 2006); this may be due to the lack of association with LTBP-1 and ECM. It is tempting to speculate that the high stress transmitted at myofibroblast adhesions is particularly efficient in activating TGF β 1 from a sufficiently organized ECM. This mechanism would provide active TGF β 1 only on demand of contractile cells to uphold myofibroblast differentiation during the contraction phase of tissue repair.

When the work is done: the fate of myofibroblasts

Solving the question whether myofibroblast differentiation is terminal or reversible will help to develop strategies to specifically and ultimately inhibit myofibroblast formation. Several studies demonstrate downregulation of α -SMA in cultured differentiated myofibroblasts in response to a variety

of factors, like TGFβ1 antagonists, growth factors, cell density, and matrix compliance without obvious signs of increased cell death. *In vivo*, however, reversal of the myofibroblast has not been documented and massive apoptosis occurs after wound healing and re-epithelialization (Desmouliere *et al.*, 1995); the signals leading to myofibroblast group suicide are still unclear. Alternatively to growth factor signals, two mechanisms are conceivable to account for myofibroblast disappearance at the end of physiological tissue repair: (a) cell release from stress and (b) increased formation of specific cell-cell contacts. In properly repaired tissue, the ECM regains its original mechanical properties and again stress-shields the resident fibroblasts. Preventing the tension release of granulation tissue by splinting the wound inhibits myofibroblast apoptosis in an animal model; subsequent splint release induces cell death (Carlson *et al.*, 2003). Similarly, apoptosis is induced by relaxing fibroblast-populated attached collagen gels induces *in vitro* (Grinnell *et al.*, 1999; Niland *et al.*, 2001). The intracellular pathways linking stress release to induction of apoptosis are unclear but likely involve cell matrix junctions (Stupack and Cheresch, 2002; Reddig and Juliano, 2005), that is to say stress perception mechanisms similar to those used to control fibroblast-to-myofibroblast transition.

Furthermore, adhesion-dependent survival signals are transmitted via cell-cell contacts and cadherins (Jamora and Fuchs, 2002; Yap and Kovacs, 2003) and formation of homotypic cadherin junctions is a possible signal of the presence of myofibroblast accumulation in late granulation tissue. Consistently, corneal myofibroblasts in dense culture significantly decrease the expression of α-SMA and de-differentiate into α-SMA-negative fibroblasts; this has been attributed to contact-induced desensitization to TGFβ1 (Petridou *et al.*, 2000). It remains to be investigated whether increased formation of OB-cadherin type junctions affects myofibroblast survival differently from N-cadherin-type contacts (Hinz *et al.*, 2004). The

potential involvement of both, specific cell-matrix and cell-cell contacts in regulating myofibroblast survival has the advantage of their extracellular accessibility to drugs.

Concluding remarks

Most efforts to understanding the molecular mechanism of myofibroblast differentiation and function are driven by the motivation to eliminate this cell in pathological tissue contractures. Although this may be difficult to accept from the sentimental point of view of someone who devotes his research to this cell, novel strategies and drugs are indeed needed to specifically counteract myofibroblast action. Inhibiting the pleiotrophic action of profibrotic cytokines such as TGFβ1 bears the disadvantage of affecting a multitude of cells that are also involved in tissue repair. **One promising antifibrosis approach is interfering with myofibroblast mechano-perception via cell-to-matrix and cell-to-cell adhesions. Just like many human individuals, myofibroblasts work best under stress; stress release leads to instant loss of internal tension – a condition that apparently drives myofibroblasts into suicide.**

CONFLICT OF INTEREST

The author states no conflict of interest.

ACKNOWLEDGMENTS

I thank Dr J.-J. Meister (Laboratory of Cell Biophysics, EPFL, Lausanne, Switzerland) for his continuous support and for providing laboratory facilities and Mrs J. Smith-Clerc for outstanding technical assistance. Dr G. Gabbiani (Department of Pathology, CMU, University of Geneva, Switzerland) is acknowledged for critically reading the paper and for his extremely helpful comments. I am grateful to Dr S. Phan (Department of Pathology, University of Michigan Medical School, MI) for his constructive contribution to the section on α-SMA transcription regulation. The work of B.H. is supported by the Swiss National Science Foundation, Grant no. 3100A0-102150/1.

REFERENCES

Abe R, Donnelly SC, Peng T, Bucala R, Metz CN (2001) Peripheral blood fibrocytes: differentiation pathway and migration to wound sites. *J Immunol* 166:7556-62

Annes JP, Chen Y, Munger JS, Rifkin DB (2004) Integrin αVβ6-mediated activation of latent TGF-β requires the latent TGF-β binding protein-1. *J Cell Biol* 165: 723-34

Annes JP, Munger JS, Rifkin DB (2003) Making sense of latent TGFβ activation. *J Cell Sci* 116:217-24

Arora PD, Narani N, McCulloch CA (1999) The compliance of collagen gels regulates transforming growth factor-beta induction of alpha-smooth muscle actin in fibroblasts. *Am J Pathol* 154:871-82

Asano Y, Ihn H, Yamane K, Jinnin M, Mimura Y, Tamaki K (2005a) Increased expression of integrin alpha(v)beta3 contributes to the establishment of autocrine TGF-beta signaling in scleroderma fibroblasts. *J Immunol* 175:7708-18

Asano Y, Ihn H, Yamane K, Jinnin M, Mimura Y, Tamaki K (2005b) Involvement of alphavbeta5 integrin-mediated activation of latent transforming growth factor beta1 in autocrine transforming growth factor beta signaling in systemic sclerosis fibroblasts. *Arthritis Rheum* 52:2897-905

Bershadsky AD, Ballestrem C, Carramusa L, Zilberman Y, Gilquin B, Khochbin S *et al.* (2006) Assembly and mechanosensory function of focal adhesions: experiments and models. *Eur J Cell Biol* 85:165-73

Bissell MJ, Radisky D (2001) Putting tumours in context. *Nat Rev Cancer* 1:46-54

Bogatkevich GS, Tourkina E, Abrams CS, Harley RA, Silver RM, Ludwicka-Bradley A (2003) Contractile activity and smooth muscle-α actin organization in thrombin-induced human lung myofibroblasts. *Am J Physiol Lung Cell Mol Physiol* 285:L334-43

Bouzeghrane F, Mercure C, Reudelhuber TL, Thibault G (2004) Alpha8beta1 integrin is upregulated in myofibroblasts of fibrotic and scarring myocardium. *J Mol Cell Cardiol* 36: 343-53

Brown RD, Ambler SK, Mitchell MD, Long CS (2005) The cardiac fibroblast: therapeutic target in myocardial remodeling and failure. *Annu Rev Pharmacol Toxicol* 45:657-87

Campaner AB, Ferreira LM, Gragnani A, Bruder JM, Cusick JL, Morgan JR (2006) Upregulation of TGF-beta1 expression may be necessary but is not sufficient for excessive scarring. *J Invest Dermatol* 126:1168-76

Carlini LE, Getz MJ, Strauch AR, Kelm RJ Jr (2002) Cryptic MCAT enhancer regulation in fibroblasts and smooth muscle cells. Suppression of tef-1 mediated activation by the single-stranded DNA-binding proteins, puralpha, purbeta, and msy1. *J Biol Chem* 277: 8682-92

Carlson MA, Longaker MT (2004) The fibroblast-populated collagen matrix as a model of wound healing: a review of the evidence. *Wound Repair Regen* 12:134-47

Carlson MA, Longaker MT, Thompson JS (2003) Wound splinting regulates granulation tissue survival. *J Surg Res* 110:304-9

Chambers RC, Leoni P, Kaminski N, Laurent GJ, Heller RA (2003) Global expression profiling of fibroblast responses to transforming growth factor-beta1 reveals the induction of inhibitor of differentiation-1 and provides evidence of smooth muscle cell phenotypic switching. *Am J Pathol* 162:533-46

- Chaponnier C, Goethals M, Janmey PA, Gabbiani F, Gabbiani G, Vandekerckhove J (1995) The specific NH2-terminal sequence Ac-EEED of alpha-smooth muscle actin plays a role in polymerization *in vitro* and *in vivo*. *J Cell Biol* 130:887–95
- Chen CS, Tan J, Tien J (2004) Mechanotransduction at cell-matrix and cell-cell contacts. *Annu Rev Biomed Eng* 6:275–302
- Chiquet-Ehrismann R, Chiquet M (2003) Tenascins: regulation and putative functions during pathological stress. *J Pathol* 200:488–99
- Christen T, Verin V, Bochaton-Piallat M, Popowski Y, Ramaekers F, Debruyne P *et al.* (2001) Mechanisms of neointima formation and remodeling in the porcine coronary artery. *Circulation* 103:882–8
- Clement S, Hinz B, Dugina V, Gabbiani G, Chaponnier C (2005) The N-terminal Ac-EEED sequence plays a role in α -smooth muscle actin incorporation into stress fibers. *J Cell Sci* 118:1395–404
- Cogan JG, Subramanian SV, Polikandriotis JA, Kelm RJ Jr, Strauch AR (2002) Vascular smooth muscle alpha-actin gene transcription during myofibroblast differentiation requires Sp1/3 protein binding proximal to the MCAT enhancer. *J Biol Chem* 277:36433–42
- De Wever O, Mareel M (2003) Role of tissue stroma in cancer cell invasion. *J Pathol* 200:429–47
- De Wever O, Nguyen QD, Van Hoorde L, Bracke M, Bruyneel E, Gespach C *et al.* (2004) Tenascin-C and SF/HGF produced by myofibroblasts *in vitro* provide convergent pro-invasive signals to human colon cancer cells through RhoA and Rac. *FASEB J* 18:1016–8
- Derynck R, Zhang YE (2003) Smad-dependent and Smad-independent pathways in TGF-beta family signalling. *Nature* 425:577–84
- Desmouliere A, Chaponnier C, Gabbiani G (2005) Tissue repair, contraction, and the myofibroblast. *Wound Repair Regen* 13:7–12
- Desmouliere A, Darby IA, Gabbiani G (2003) Normal and pathologic soft tissue remodeling: role of the myofibroblast, with special emphasis on liver and kidney fibrosis. *Lab Invest* 83:1689–707
- Desmouliere A, Geinoz A, Gabbiani F, Gabbiani G (1993) Transforming growth factor-beta 1 induces alpha-smooth muscle actin expression in granulation tissue myofibroblasts and in quiescent and growing cultured fibroblasts. *J Cell Biol* 122:103–11
- Desmouliere A, Guyot C, Gabbiani G (2004) The stroma reaction myofibroblast: a key player in the control of tumor cell behavior. *Int J Dev Biol* 48:509–17
- Desmouliere A, Redard M, Darby I, Gabbiani G (1995) Apoptosis mediates the decrease in cellularity during the transition between granulation tissue and scar. *Am J Pathol* 146:56–66
- Direkze NC, Forbes SJ, Brittan M, Hunt T, Jeffery R, Preston SL *et al.* (2003) Multiple organ engraftment by bone-marrow-derived myofibroblasts and fibroblasts in bone-marrow-transplanted mice. *Stem Cells* 21:514–20
- Discher DE, Janmey P, Wang YL (2005) Tissue cells feel and respond to the stiffness of their substrate. *Science* 310:1139–43
- Dooley S, Said HM, Gressner AM, Floege J, En-Nia A, Mertens PR (2006) Y-box protein-1 is the crucial mediator of antifibrotic interferon-gamma effects. *J Biol Chem* 281:1784–95
- Dugina V, Fontao L, Chaponnier C, Vasiliev J, Gabbiani G (2001) Focal adhesion features during myofibroblastic differentiation are controlled by intracellular and extracellular factors. *J Cell Sci* 114:3285–96
- Eckes B, Krieg T (2004) Regulation of connective tissue homeostasis in the skin by mechanical forces. *Clin Exp Rheumatol* 22:S73–6
- Evans RA, Tian YC, Steadman R, Phillips AO (2003) TGF-beta1-mediated fibroblast-myofibroblast terminal differentiation—the role of Smad proteins. *Exp Cell Res* 282:90–100
- Eyden B (2005) The myofibroblast: a study of normal, reactive and neoplastic tissues, with an emphasis on ultrastructure. Part 1 – normal and reactive cells. *J Submicroscop Cytol Pathol* 37:109–204
- Faouzi S, Le Bail B, Neaud V, Boussarie L, Saric J, Bioulac-Sage P *et al.* (1999) Myofibroblasts are responsible for collagen synthesis in the stroma of human hepatocellular carcinoma: an *in vivo* and *in vitro* study. *J Hepatol* 30:275–84
- Feng XH, Derynck R (2005) Specificity and versatility in TGF-beta signaling through Smads. *Annu Rev Cell Dev Biol* 21:659–93
- Folger PA, Zekaria D, Grotendorst G, Masur SK (2001) Transforming growth factor-beta-stimulated connective tissue growth factor expression during corneal myofibroblast differentiation. *Invest Ophthalmol Vis Sci* 42:2534–41
- Forbes SJ, Russo FP, Rey V, Burra P, Rugge M, Wright NA *et al.* (2004) A significant proportion of myofibroblasts are of bone marrow origin in human liver fibrosis. *Gastroenterology* 126:955–63
- Fung YC (1993) *Mechanical properties of living tissues*. 2nd ed. New York, NY: Springer Verlag
- Gabbiani G (2003) The myofibroblast in wound healing and fibrocontractive diseases. *J Pathol* 200:500–3
- Gallucci RM, Lee EG, Tomasek JJ (2006) IL-6 modulates alpha-smooth muscle actin expression in dermal fibroblasts from IL-6-deficient mice. *J Invest Dermatol* 126:561–8
- Geiger B, Bershadsky A, Pankov R, Yamada KM (2001) Transmembrane crosstalk between the extracellular matrix – cytoskeleton crosstalk. *Nat Rev Mol Cell Biol* 2:793–805
- Glukhova MA, Frid MG, Shekhnin BV, Vasilevska TD, Grunwald J, Saginati M *et al.* (1989) Expression of extra domain A fibronectin sequence in vascular smooth muscle cells is phenotype dependent. *J Cell Biol* 109:357–66
- Goffin JM, Pittet P, Csucs G, Lussi JW, Meister JJ, Hinz B (2006) Focal adhesion size controls tension-dependent recruitment of alpha-smooth muscle actin to stress fibers. *J Cell Biol* 172:259–68
- Greenberg RS, Bernstein AM, Benezra M, Gelman IH, Taliana L, Masur SK (2006) FAK-dependent regulation of myofibroblast differentiation. *FASEB J* 20:1006–8
- Greenleaf JF, Fatemi M, Insana M (2003) Selected methods for imaging elastic properties of biological tissues. *Annu Rev Biomed Eng* 5:57–78
- Grinnell F (2003) Fibroblast biology in three-dimensional collagen matrices. *Trends Cell Biol* 13:264–9
- Grinnell F, Ho CH (2002) Transforming growth factor beta stimulates fibroblast-collagen matrix contraction by different mechanisms in mechanically loaded and unloaded matrices. *Exp Cell Res* 273:248–55
- Grinnell F, Zhu M, Carlson MA, Abrams JM (1999) Release of mechanical tension triggers apoptosis of human fibroblasts in a model of regressing granulation tissue. *Exp Cell Res* 248:608–19
- Groma V (1998) Demonstration of collagen type VI and alpha-smooth muscle actin in renal fibrotic injury in man. *Nephrol Dial Transplant* 13:305–12
- Gross J, Farinelli W, Sadow P, Anderson R, Bruns R (1995) On the mechanism of skin wound “contraction”: a granulation tissue “knock-out” with a normal phenotype. *Proc Natl Acad Sci USA* 92:5982–6
- Grotendorst GR, Duncan MR (2005) Individual domains of connective tissue growth factor regulate fibroblast proliferation and myofibroblast differentiation. *FASEB J* 19:729–38
- Gumbiner BM (2005) Regulation of cadherin-mediated adhesion in morphogenesis. *Nat Rev Mol Cell Biol* 6:622–34
- Guyot C, Lepreux S, Combe C, Doudnikoff E, Bioulac-Sage P, Balabaud C *et al.* (2006) Hepatic fibrosis and cirrhosis: the (myo)fibroblastic cell subpopulations involved. *Int J Biochem Cell Biol* 38:135–51
- Hagood JS, Prabhakaran P, Kumbla P, Salazar L, MacEwen MW, Barker TH *et al.* (2005) Loss of fibroblast Thy-1 expression correlates with lung fibrogenesis. *Am J Pathol* 167:365–79
- Hashimoto N, Jin H, Liu T, Chensue SW, Phan SH (2004) Bone marrow-derived progenitor cells in pulmonary fibrosis. *J Clin Invest* 113:243–52
- Hautmann MB, Madsen CS, Owens GK (1997) A transforming growth factor beta (TGFbeta) control element drives TGFbeta-induced stimulation of smooth muscle alpha-actin gene expression in concert with two CArG elements. *J Biol Chem* 272:10948–56
- Henderson NC, Mackinnon AC, Farnworth SL, Poirier F, Russo FP, Iredale JP *et al.* (2006) Galectin-3 regulates myofibroblast activation and hepatic fibrosis. *Proc Natl Acad Sci USA* 103:5060–5
- Higashi K, Inagaki Y, Fujimori K, Nakao A, Kaneko H, Nakatsuka I (2003) Interferon-gamma interferes with transforming growth factor-beta signaling through direct interac-

- tion of YB-1 with Smad3. *J Biol Chem* 278:43470-9
- Hinz B (2006) Masters and servants of the force: the role of matrix adhesions in myofibroblast force perception and transmission. *Eur J Cell Biol* 85:175-81
- Hinz B, Celetta G, Tomasek JJ, Gabbiani G, Chaponnier C (2001a) Alpha-smooth muscle actin expression upregulates fibroblast contractile activity. *Mol Biol Cell* 12:2730-41
- Hinz B, Dugina V, Ballestrem C, Wehrle-Haller B, Chaponnier C (2003) Alpha-smooth muscle actin is crucial for focal adhesion maturation in myofibroblasts. *Mol Biol Cell* 14:2508-19
- Hinz B, Gabbiani G (2003a) Cell-matrix and cell-cell contacts of myofibroblasts: role in connective tissue remodeling. *Thromb Haemost* 90:993-1002
- Hinz B, Gabbiani G (2003b) Mechanisms of force generation and transmission by myofibroblasts. *Curr Opin Biotechnol* 14:538-46
- Hinz B, Gabbiani G, Chaponnier C (2002) The NH2-terminal peptide of alpha-smooth muscle actin inhibits force generation by the myofibroblast *in vitro* and *in vivo*. *J Cell Biol* 157:657-63
- Hinz B, Mastrangelo D, Iselin CE, Chaponnier C, Gabbiani G (2001b) Mechanical tension controls granulation tissue contractile activity and myofibroblast differentiation. *Am J Pathol* 159:1009-20
- Hinz B, Pittet P, Smith-Clerc J, Chaponnier C, Meister JJ (2004) Myofibroblast development is characterized by specific cell-cell adhesion junctions. *Mol Biol Cell* 15:4310-20
- Hu B, Wu Z, Phan SH (2003) Smad3 mediates transforming growth factor-beta-induced alpha-smooth muscle actin expression. *Am J Respir Cell Mol Biol* 29:397-404
- Ishii G, Sangai T, Oda T, Aoyagi Y, Hasebe T, Kanomata N *et al.* (2003) Bone-marrow-derived myofibroblasts contribute to the cancer-induced stromal reaction. *Biochem Biophys Res Commun* 309:232-40
- Ishii G, Sangai T, Sugiyama K, Ito T, Hasebe T, Endoh Y *et al.* (2005) *In vivo* characterization of bone marrow-derived fibroblasts recruited into fibrotic lesions. *Stem Cells* 23:699-706
- Jamora C, Fuchs E (2002) Intercellular adhesion, signalling and the cytoskeleton. *Nat Cell Biol* 4:E101-8
- Kanangat S, Postlethwaite AE, Higgins GC, Hasty KA (2006) Novel functions of intracellular IL-1ra in human dermal fibroblasts: implications in the pathogenesis of fibrosis. *J Invest Dermatol* 126:756-65
- Katsumi A, Orr AW, Tzima E, Schwartz MA (2004) Integrins in mechanotransduction. *J Biol Chem* 279:12001-4
- Kaufman LJ, Brangwynne CP, Kasza KE, Filippidi E, Gordon VD, Deisboeck TS *et al.* (2005) Glioma expansion in collagen I matrices: analyzing collagen concentration-dependent growth and motility patterns. *Biophys J* 89:635-50
- Keski-Oja J, Koli K, von Melchner H (2004) TGF-beta activation by traction? *Trends Cell Biol* 14:657-9
- Kitamura M, Shimizu M, Ino H, Okeie K, Yamaguchi M, Funjio N *et al.* (2001) Collagen remodeling and cardiac dysfunction in patients with hypertrophic cardiomyopathy: the significance of type III and VI collagens. *Clin Cardiol* 24:325-9
- Ko KS, Arora PD, McCulloch CA (2001) Cadherins mediate intercellular mechanical signaling in fibroblasts by activation of stretch-sensitive calcium-permeable channels. *J Biol Chem* 276:35967-77
- Koli K, Hyytiainen M, Ryyanen MJ, Keski-Oja J (2005) Sequential deposition of latent TGF-beta binding proteins (LTBPs) during formation of the extracellular matrix in human lung fibroblasts. *Exp Cell Res* 310:370-82
- Koumas L, Smith TJ, Feldon S, Blumberg N, Phipps RP (2003) Thy-1 expression in human fibroblast subsets defines myofibroblastic or lipofibroblastic phenotypes. *Am J Pathol* 163:1291-300
- Lan HY (2003) Tubular epithelial-myofibroblast transdifferentiation mechanisms in proximal tubule cells. *Curr Opin Nephrol Hypertens* 12:25-9
- Leask A, Holmes A, Abraham DJ (2002) Connective tissue growth factor: a new and important player in the pathogenesis of fibrosis. *Curr Rheumatol Rep* 4:136-42
- Levine D, Rockey DC, Milner TA, Breuss JM, Fallon JT, Schnapp LM (2000) Expression of the integrin alpha8beta1 during pulmonary and hepatic fibrosis. *Am J Pathol* 156:1927-35
- Liotta LA, Kohn EC (2001) The microenvironment of the tumour-host interface. *Nature* 411:375-9
- Liu C, Gaca MD, Swenson ES, Vellucci VF, Reiss M, Wells RG (2003) Smads 2 and 3 are differentially activated by transforming growth factor-beta (TGF-beta) in quiescent and activated hepatic stellate cells. Constitutive nuclear localization of Smads in activated cells is TGF-beta-independent. *J Biol Chem* 278:11721-8
- Liu T, Dhanasekaran SM, Jin H, Hu B, Tomlins SA, Chinnaiyan AM *et al.* (2004) FIZ1 stimulation of myofibroblast differentiation. *Am J Pathol* 164:1315-26
- Lu M, Munger JS, Steadele M, Busald C, Tellier M, Schnapp LM (2002) Integrin alpha8beta1 mediates adhesion to LAP-TGFbeta1. *J Cell Sci* 115:4641-8
- Ludbrook SB, Barry ST, Delves CJ, Horgan CM (2003) The integrin alphavbeta3 is a receptor for the latency-associated peptides of transforming growth factors beta1 and beta3. *Biochem J* 369:311-8
- Malmstrom J, Lindberg H, Lindberg C, Bratt C, Wieslander E, Delander EL *et al.* (2004) Transforming growth factor-beta 1 specifically induce proteins involved in the myofibroblast contractile apparatus. *Mol Cell Proteomics* 3:466-77
- Mangasser-Stephan K, Gartung C, Lahme B, Gressner AM (2001) Expression of isoforms and splice variants of the latent transforming growth factor beta binding protein (LTBP) in cultured human liver myofibroblasts. *Liver* 21:105-13
- Marenzana M, Wilson-Jones N, Mudera V, Brown RA (2006) The origins and regulation of tissue tension: identification of collagen tension-fixation process *in vitro*. *Exp Cell Res* 312:423-33
- Martinac B (2004) Mechanosensitive ion channels: molecules of mechanotransduction. *J Cell Sci* 117:2449-60
- Massague J, Seoane J, Wotton D (2005) Smad transcription factors. *Genes Dev* 19:2783-810
- Micera A, Vigneti E, Pickholtz D, Reich R, Pappo O, Bonini S *et al.* (2001) Nerve growth factor displays stimulatory effects on human skin and lung fibroblasts, demonstrating a direct role for this factor in tissue repair. *Proc Natl Acad Sci USA* 98:6162-7
- Mimura Y, Ihn H, Jinnin M, Asano Y, Yamane K, Tamaki K (2005) Constitutive phosphorylation of focal adhesion kinase is involved in the myofibroblast differentiation of scleroderma fibroblasts. *J Invest Dermatol* 124:886-92
- Mori L, Bellini A, Stacey MA, Schmidt M, Mattoli S (2005) Fibrocytes contribute to the myofibroblast population in wounded skin and originate from the bone marrow. *Exp Cell Res* 304:81-90
- Moustakas A, Heldin CH (2005) Non-Smad TGF-beta signals. *J Cell Sci* 118:3573-84
- Mueller MM, Fusenig NE (2004) Friends or foes - bipolar effects of the tumour stroma in cancer. *Nat Rev Cancer* 4:839-49
- Munger JS, Huang X, Kawakatsu H, Griffiths MJ, Dalton SL, Wu J *et al.* (1999) The integrin alpha v beta 6 binds and activates latent TGF beta 1: a mechanism for regulating pulmonary inflammation and fibrosis. *Cell* 96:319-28
- Mykkanen OM, Gronholm M, Ronty M, Lalowski M, Salmikangas P, Suila H *et al.* (2001) Characterization of human palladin, a microfilament-associated protein. *Mol Biol Cell* 12:3060-73
- Naugle JE, Olson ER, Zhang X, Mase SE, Pilati CF, Maron MB *et al.* (2006) Type VI collagen induces cardiac myofibroblast differentiation: implications for postinfarction remodeling. *Am J Physiol Heart Circ Physiol* 290:H323-30
- Ng CP, Hinz B, Swartz MA (2005) Interstitial fluid flow induces myofibroblast differentiation and collagen alignment *in vitro*. *J Cell Sci* 118:4731-9
- Niland S, Cremer A, Fluck J, Eble JA, Krieg T, Sollberg S (2001) Contraction-dependent apoptosis of normal dermal fibroblasts. *J Invest Dermatol* 116:686-92
- Paszek MJ, Zahir N, Johnson KR, Lakins JN, Rozenberg GI, Gefen A *et al.* (2005) Tensional homeostasis and the malignant phenotype. *Cancer Cell* 8:241-54
- Petridou S, Maltseva O, Spanakis S, Masur SK (2000) TGF-beta receptor expression and smad2 localization are cell density dependent

- dent in fibroblasts. *Invest Ophthalmol Vis Sci* 41:89-95
- Phan SH (2002) The myofibroblast in pulmonary fibrosis. *Chest* 122:286S-95S
- Rajkumar VS, Howell K, Csiszar K, Denton CP, Black CM, Abraham DJ (2005) Shared expression of phenotypic markers in systemic sclerosis indicates a convergence of pericytes and fibroblasts to a myofibroblast lineage in fibrosis. *Arthritis Res Ther* 7:R1113-23
- Reddig PJ, Juliano RL (2005) Clinging to life: cell to matrix adhesion and cell survival. *Cancer Metastasis Rev* 24:425-39
- Ronnov-Jessen L, Petersen OW (1993) Induction of alpha-smooth muscle actin by transforming growth factor-beta 1 in quiescent human breast gland fibroblasts. Implications for myofibroblast generation in breast neoplasia. *Lab Invest* 68:696-707
- Rönty M, Leivonen S-K, Hinz B, Rachlin A, Otey C, Kähäri V-M *et al.* (2006) Isoform-specific regulation of the actin-organizing protein plectin during TGF-beta1-induced myofibroblast differentiation. *J Invest Dermatol* 126:2387-96
- Roy SG, Nozaki Y, Phan SH (2001) Regulation of alpha-smooth muscle actin gene expression in myofibroblast differentiation from rat lung fibroblasts. *Int J Biochem Cell Biol* 33:723-34
- Sartore S, Chiavegato A, Faggin E, Franch R, Puato M, Ausoni S *et al.* (2001) Contribution of adventitial fibroblasts to neointima formation and vascular remodeling: from innocent bystander to active participant. *Circ Res* 89:1111-21
- Schildmeyer LA, Braun R, Taffet G, Debiase M, Burns AE, Bradley A *et al.* (2000) Impaired vascular contractility and blood pressure homeostasis in the smooth muscle alpha-actin null mouse. *FASEB J* 14:2213-20
- Schnabl B, Kweon YO, Frederick JP, Wang XF, Rippe RA, Brenner DA (2001) The role of Smad3 in mediating mouse hepatic stellate cell activation. *Hepatology* 34:89-100
- Schurch W, Seemayer TA, Hinz B, Gabbiani G (2006) The myofibroblast. In: *Histology for pathologists* (Mills SE, ed), Philadelphia, USA: Lippincott-Williams & Wilkins. Pub. (in press)
- Serini G, Bochaton-Piallat ML, Ropraz P, Geinoz A, Borsi L, Zardi L *et al.* (1998) The fibronectin domain ED-A is crucial for myofibroblastic phenotype induction by transforming growth factor-beta1. *J Cell Biol* 142:873-81
- Shephard P, Hinz B, Smola-Hess S, Meister JJ, Krieg T, Smola H (2004a) Dissecting the roles of endothelin, TGF-beta and GM-CSF on myofibroblast differentiation by keratinocytes. *Thromb Haemostasis* 92:262-74
- Shephard P, Martin G, Smola-Hess S, Brunner G, Krieg T, Smola H (2004b) Myofibroblast differentiation is induced in keratinocyte-fibroblast co-cultures and is antagonistically regulated by endogenous transforming growth factor-beta and interleukin-1. *Am J Pathol* 164:2055-66
- Shi-Wen X, Chen Y, Denton CP, Eastwood M, Renzoni EA, Bou-Gharios G *et al.* (2004) Endothelin-1 promotes myofibroblast induction through the ETA receptor via a rac1-phosphoinositide 3-kinase/Akt-dependent pathway and is essential for the enhanced contractile phenotype of fibrotic fibroblasts. *Mol Biol Cell* 15:2707-19
- Stupack DG, Cheresch DA (2002) Get a ligand, get a life: integrins, signaling and cell survival. *J Cell Sci* 115:3729-38
- Subramanian SV, Polikandriotis JA, Kelm RJ Jr, David JJ, Orosz CG, Strauch AR (2004) Induction of vascular smooth muscle alpha-actin gene transcription in transforming growth factor beta1-activated myofibroblasts mediated by dynamic interplay between the Pur repressor proteins and Sp1/Smad coactivators. *Mol Biol Cell* 15:4532-43
- Swaney JS, Roth DM, Olson ER, Naugle JE, Meszaros JG, Insel PA (2005) Inhibition of cardiac myofibroblast formation and collagen synthesis by activation and overexpression of adenylyl cyclase. *Proc Natl Acad Sci USA* 102:437-42
- Swartz EA, Johnson AD, Owens GK (1998) Two MCAT elements of the SM alpha-actin promoter function differentially in SM vs non-SM cells. *Am J Physiol* 275:C608-18
- Tamaoki M, Imanaka-Yoshida K, Yokoyama K, Nishioka T, Inada H, Hiroe M *et al.* (2005) Tenascin-C regulates recruitment of myofibroblasts during tissue repair after myocardial injury. *Am J Pathol* 167:71-80
- Tamariz E, Grinnell F (2002) Modulation of fibroblast morphology and adhesion during collagen matrix remodeling. *Mol Biol Cell* 13:3915-29
- Thannickal VJ, Lee DY, White ES, Cui Z, Larios JM, Chacon R *et al.* (2003) Myofibroblast differentiation by transforming growth factor-beta1 is dependent on cell adhesion and integrin signaling via focal adhesion kinase. *J Biol Chem* 278:12384-9
- Thannickal VJ, Toews GB, White ES, Lynch JP III, Martinez FJ (2004) Mechanisms of pulmonary fibrosis. *Annu Rev Med* 55:395-417
- Tomasek JJ, Gabbiani G, Hinz B, Chaponnier C, Brown RA (2002) Myofibroblasts and mechano-regulation of connective tissue remodeling. *Nat Rev Mol Cell Biol* 3:349-63
- Tomasek JJ, McRae J, Owens GK, Haaksma CJ (2005) Regulation of alpha-smooth muscle actin expression in granulation tissue myofibroblasts is dependent on the intronic CARG element and the transforming growth factor-beta1 control element. *Am J Pathol* 166:1343-51
- Tomita K, van Bokhoven A, van Leenders GJ, Ruijter ET, Jansen CF, Bussemakers MJ *et al.* (2000) Cadherin switching in human prostate cancer progression. *Cancer Res* 60:3650-4
- Uemura M, Swenson ES, Gaca MD, Giordano FJ, Reiss M, Wells RG (2005) Smad2 and Smad3 play different roles in rat hepatic stellate cell function and alpha-smooth muscle actin organization. *Mol Biol Cell* 16:4214-24
- Unsold C, Hyytiäinen M, Bruckner-Tuderman L, Keski-Oja J (2001) Latent TGF-beta binding protein LTBP-1 contains three potential extracellular matrix interacting domains. *J Cell Sci* 114:187-97
- van der Loop FT, Schaart G, Timmer ED, Ramaekers FC, van Eys GJ (1996) Smoothelin, a novel cytoskeletal protein specific for smooth muscle cells. *J Cell Biol* 134:401-11
- Virag JJ, Murry CE (2003) Myofibroblast and endothelial cell proliferation during murine myocardial infarct repair. *Am J Pathol* 163:2433-40
- Vogel V, Sheetz M (2006) Local force and geometry sensing regulate cell functions. *Nat Rev Mol Cell Biol* 7:265-75
- Wang J, Zohar R, McCulloch CA (2006) Multiple roles of alpha-smooth muscle actin in mechanotransduction. *Exp Cell Res* 312:205-14
- Wells RG (2005) The role of matrix stiffness in hepatic stellate cell activation and liver fibrosis. *J Clin Gastroenterol* 39:S158-61
- Werner S, Grose R (2003) Regulation of wound healing by growth factors and cytokines. *Physiol Rev* 83:835-70
- Wilkes MC, Mitchell H, Penheiter SG, Dore JJ, Suzuki K, Edens M *et al.* (2005) Transforming growth factor-beta activation of phosphatidylinositol 3-kinase is independent of Smad2 and Smad3 and regulates fibroblast responses via p21-activated kinase-2. *Cancer Res* 65:10431-40
- Yang L, Qiu CX, Ludlow A, Ferguson MW, Brunner G (1999) Active transforming growth factor-beta in wound repair: determination using a new assay. *Am J Pathol* 154:105-11
- Yap AS, Kovacs EM (2003) Direct cadherin-activated cell signaling: a view from the plasma membrane. *J Cell Biol* 160:11-6
- Yeung T, Georges PC, Flanagan LA, Marg B, Ortiz M, Funaki M *et al.* (2005) Effects of substrate stiffness on cell morphology, cytoskeletal structure, and adhesion. *Cell Motil Cytoskeleton* 60:24-34
- Zalewski A, Shi Y, Johnson AG (2002) Diverse origin of intimal cells: smooth muscle cells, myofibroblasts, fibroblasts, and beyond? *Circ Res* 91:652-5
- Zhou Y, Hagood JS, Murphy-Ullrich JE (2004) Thy-1 expression regulates the ability of rat lung fibroblasts to activate transforming growth factor-beta in response to fibrogenic stimuli. *Am J Pathol* 165:659-69